

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date,
3 February 2005 (03.02.2005)

PCT

(10) International Publication Number
WO 2005/009418 A2

- (51) International Patent Classification⁷: **A61K 31/00**
- (21) International Application Number:
PCT/TB2004/002703
- (22) International Filing Date: 26 July 2004 (26.07.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/490,216 25 July 2003 (25.07.2003) US
- (71) Applicants (for all designated States except US): **IDENIX (CAYMAN) LIMITED**; Walker Secretaries, Walker House, Grand Cayman (KY). **CENTRE NATIONAL DE LA RECHERCHE SCIENTIFIQUE** [FR/FR]; 3, rue Michel-Ange, F-75794 Paris Cédex 16 (FR). **L'UNIVERSITE MONTPELLIER II** [FR/FR]; 2 Place Eugène Bataillon, F-34095 Montpellier Cedex 5 (FR).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **STORER, Richard** [GB/GB]; Sandgate Point, The Leas, Folkestone, Kent CT20 2JE (GB). **GOSSELIN, Gilles** [FR/FR]; 487, rue Celleneuve St. Hilaire, F-34070 Montpellier (FR). **DUKHAN, David** [FR/FR]; 42 Impasse Tour Buffel, Res. les Verriers, F-34070 Montpellier (FR). **LEROY, Frédéric** [FR/FR]; Rue de la Font du Loupiou, F-34725 Jonquières (FR).
- (74) Agents: **MARSHALL, Cameron, John et al.**; Carpmeals & Ransford, 43-45 Bloomsbury Square, London WC1A 2RA (GB).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PURIN NUCLEOSIDE ANALOGUES FOR TREATING FLAVIVIRIDAE INCLUDING HEPATITIS C

(57) Abstract: This invention is directed to a method for treating a host, especially a human, infected with hepatitis C, flavivirus and/or pestivirus, comprising administering to that host an effective amount of an anti-HCV biologically active pentofuranonucleoside where the pentofuranonucleoside base is an optionally substituted 2-azapurine. The optionally substituted pentofuranonucleoside, or a salt or prodrug thereof, may be administered alone or in combination with one or more optionally substituted pentofuranonucleosides or other anti-viral agents.



WO 2005/009418 A2

PURINE NUCLEOSIDE ANALOGUES FOR TREATING *FLAVIVIRIDAE* INCLUDING HEPATITIS C

This application claims priority to U.S. Provisional Application No. 60/490,216,
5 filed July 25, 2003.

FIELD OF THE INVENTION

This invention is in the area of pharmaceutical chemistry, and, in particular, in the area of purine nucleosides, their syntheses, and their use as anti-*Flaviviridae* agents in the treatment of hosts infected with *Flaviviridae* and especially with Hepatitis C.

BACKGROUND OF THE INVENTION

Flaviviridae Viruses

The *Flaviviridae* family of viruses comprises at least three distinct genera: *pestiviruses*, which cause disease in cattle and pigs; *flaviviruses*, which are the primary cause of diseases such as dengue fever and yellow fever; and *hepaciviruses* such as hepatitis C (HCV). The flavivirus genus includes more than 68 members separated into groups on the basis of serological relatedness (Calisher et al., *J. Gen. Virol.*, 1993, 70, 37-43). Clinical symptoms vary and include fever, encephalitis and hemorrhagic fever (*Fields Virology*, Editors: Fields, B. N., Knipe, D. M., and Howley, P. M., Lippincott-Raven Publishers, Philadelphia, PA, 1996, Chapter 31, 931-959). Flaviviruses of global concern that are associated with human disease include Dengue virus, hemorrhagic fever viruses such as Lassa, Ebola, and yellow fever virus, shock syndrome, and Japanese encephalitis virus (Halstead, S. B., *Rev. Infect. Dis.*, 1984, 6, 251-264; Halstead, S. B., *Science*, 239:476-481, 1988; Monath, T. P., *New Eng. J. Med.*, 1988, 319, 641-643).

The pestivirus genus includes bovine viral diarrhea virus (BVDV), classical swine fever virus (CSFV, also called hog cholera virus) and border disease virus (BDV) of sheep (Moennig, V. et al. *Adv. Vir. Res.* 1992, 41, 53-98). Pestivirus infections of domesticated livestock (cattle, pigs and sheep) cause significant economic losses worldwide. BVDV causes mucosal disease in cattle and is of significant economic importance to the livestock industry (Meyers, G. and Thiel, H.-J., *Advances in Virus Research*, 1996, 47, 53-118; Moennig V., et al, *Adv. Vir. Res.* 1992, 41, 53-98). Human pestiviruses have not been as extensively characterized as the animal pestiviruses. However, serological surveys indicate considerable pestivirus exposure in humans.

Pestiviruses and *hepaciviruses* are closely related virus groups within the *Flaviviridae* family. Other closely related viruses in this family include the GB virus A, GB virus A-like agents, GB virus-B and GB virus-C (also called hepatitis G virus, HGV). The hepacivirus group (hepatitis C virus; HCV) consists of a number of closely related but genotypically distinguishable viruses that infect humans. There are approximately 6 HCV genotypes and more than 50 subtypes. Due to the similarities between pestiviruses and hepaciviruses, combined with the poor ability of hepaciviruses to grow efficiently in cell culture, bovine viral diarrhea virus (BVDV) is often used as a surrogate to study the HCV virus.

The genetic organization of pestiviruses and hepaciviruses is very similar. These positive stranded RNA viruses possess a single large open reading frame (ORF) encoding all the viral proteins necessary for virus replication. These proteins are expressed as a polyprotein that is co- and post-translationally processed by both cellular and virus-encoded proteinases to yield the mature viral proteins. The viral proteins responsible for the replication of the viral genome RNA are located within approximately the carboxy-terminal. Two-thirds of the ORF are termed nonstructural (NS) proteins. The genetic organization and polyprotein processing of the nonstructural protein portion of the ORF for pestiviruses and hepaciviruses is very similar. For both the pestiviruses and hepaciviruses, the mature nonstructural (NS) proteins, in sequential order from the amino-terminus of the nonstructural protein coding region to the carboxy-terminus of the ORF, consist of p7, NS1, NS2A, NS2B, NS3, NS4A, NS4B, NS5A, and NS5B.

The NS proteins of pestiviruses and hepaciviruses share sequence domains that are characteristic of specific protein functions. For example, the NS1 glycoprotein is a cell-surface protein that is translocated into the ER lumen. NS1 was characterized initially as soluble complement-fixing antigen found in sera and tissues of infected animals, and now is known to elicit humoral immune responses in its extracellular form. Antibodies to NS1 may be used to confer passive immunity to certain pestiviruses and flaviviruses. NS1 has been implicated in the process of RNA replication where it is believed to have a functional role in the cytoplasmic processing of RNA. NS2A is a small (approximately 22 kd) protein of unknown function. Studies suggest that it binds to NS3 and NS5, and so may be a recruiter of RNA templates to membrane-bound replicase. NS2B also is a small (about 14 kd) protein that is membrane-associated, and is a required cofactor for the serine protease function of NS3, with which it forms a complex.

The NS3 proteins of viruses in both groups are large (about 70 kd), membrane-associated proteins that possess amino acid sequence motifs characteristic of serine proteinases and of helicases (Gorbalenya et al. (1988) *Nature* 333:22; Bazan and Fletterick (1989) *Virology* 171:637-639; Gorbalenya et al. (1989) *Nucleic Acid Res.* 17:3889-3897). Thus, the NS3 proteins have enzymatic activity needed for processing polyproteins for RNA replication. The C-terminal end of the NS3 proteins have an RNA triphosphatase activity that appears to modify the 5' end of the genome prior to 5'-cap addition by guanylyltransferase.

NS4A and NS4B are membrane-associated, small (about 16 kd and about 27 kd, respectively), hydrophobic proteins that appear to function in RNA replication by anchoring replicase components to cellular membranes (Fields, *Virology*, 4th Edition, 2001, p. 1001).

The NS5 proteins are the largest (about 103 kd) and most conserved, with sequence homology to other (+)-stranded RNA viruses. It also plays a pivotal role in viral replication. The NS5B proteins of pestiviruses and hepaciviruses are the enzymes necessary for synthesis of the negative-stranded RNA intermediate that is complementary to the viral genome, and of the positive-stranded RNA that is complementary to the negative-stranded RNA intermediate. The NS5B gene product has Gly-Asp-Asp (GDD) as a hallmark sequence, which it shares with reverse transcriptases

and other viral polymerases and which is predictive of RNA dependent RNA polymerase (RdRP) activity (DeFrancesco et al., *Antiviral Research*, 2003, 58:1-16). Interestingly, it was found that the NS5B C-terminal 21 residue long hydrophobic tail is needed to target NS5B to the ER membrane, but its removal has no other effect and, in fact, leads to increased enzymatic solubility and activity (Tomei et al., *J. Gen. Virol.*, 2000, 81:759-767; Lohmann et al., *J. Virol.*, 1997, 71:8416-28; Ferrari et al., *J. Virol.*, 1999, 73:1649-54).

The NS5B enzyme products have the motifs characteristic of RNA-directed RNA polymerases, and in addition, share homology with methyltransferase enzymes that are involved in RNA cap formation (Koonin, E.V. and Dolja, V.V. (1993) *Crit. Rev. Biochem. Molec. Biol.* 28:375-430; Behrens et al.(1996) *EMBO J.* 15:12-22; Lohmann et al.(1997) *J. Virol.* 71:8416-8428; Yuan et al.(1997) *Biochem. Biophys. Res. Comm.* 232:231-235; Hagedorn, PCT WO 97/12033; Zhong et al.(1998) *J. Virol.* 72:9365-9369). The unliganded crystal structure of NS5B shows the unique structural feature of folding in a classic “right hand” shape, in which fingers, palm and thumb subdomains can be recognized (a feature it shares with other polymerases), but differs from other “half-open right hand” polymerases by having a more compact shapes due to two extended loops that span the finger and thumb domains at the top of the active site cavity (DeFrancesco et al. at 9). The finger, thumb and palm subdomains encircle the active site cavity to which the RNA template and NTP substrates have access via two positively charged tunnels (Bressanelli et al., *J. Virol.*, 2002, 76, 3482-92). Finger and thumb domains have strong interactions that limit their ability to change conformation independently of one another, a structural feature shared by other RdRPs. The thumb domain contains a β -hairpin loop that extends toward the cleft of the active site and may play a role in restricting the binding of the template/primer at the enzyme active site (DeFrancesco et al., at 10). Studies are in progress to determine the role of this loop in the initiation mechanism of RNA synthesis (Id.)

Nucleotidyl transfer reaction residues are located in the palm domain and contain the signature GDD motif (DeFrancesco et al., at 9). Palm domain geometry is highly conserved in all polymerases, and has a conserved two-metal-ion catalytic center that is required for catalyzing a phosphory transfer reaction at the polymerase active site.

It is believed that the de novo initiation model of RNA polymerization, rather than a "copy back" mechanism, is utilized by pesti-, flavi- and hepaciviruses. In the de novo initiation model, complementary RNA synthesis is initiated at the 3'-end of the genome by a nucleotide triphosphate rather than a nucleic acid or a protein primer.

5 Purified NS5B is capable of this type of primer-independent action, and the C-terminal β -loop is believed to correctly position the 3'-end of the RNA template by functioning as a gate that retards slippage of the RNA 3'-end through the polymerase active site (Hong et al., *Virology*, 2001, 285:6-11. Bressanelli et al. reported the structure of NS5B polymerase in complex with nucleotides in which three distinct nucleotide-binding sites

10 were observed in the catalytic center of the HCV RdRP, and the complex exhibited a geometry similar to the de novo initiation complex of phi 6 polymerase (Bressanelli et al., *J. Virol.*, 2002, 76: 3482-92). Thus, de novo initiation occurs and apparently is followed by RNA elongation, termination of polymerization, and release of the new strand. At each of these steps is the opportunity for intervention and inhibition of the

15 viral lifecycle.

The actual roles and functions of the NS proteins of pestiviruses and hepaciviruses in the lifecycle of the viruses are directly analogous. In both cases, the NS3 serine proteinase is responsible for all proteolytic processing of polyprotein precursors downstream of its position in the ORF (Wiskerchen and Collett (1991)

20 *Virology* 184:341-350; Bartenschlager et al. (1993) *J. Virol.* 67:3835-3844; Eckart et al. (1993) *Biochem. Biophys. Res. Comm.* 192:399-406; Grakoui et al. (1993) *J. Virol.* 67:2832-2843; Grakoui et al. (1993) *Proc. Natl. Acad. Sci. USA* 90:10583-10587; Hijikata et al. (1993) *J. Virol.* 67:4665-4675; Tome et al. (1993) *J. Virol.* 67:4017-4026). The NS4A protein, in both cases, acts as a cofactor with the NS3 serine protease

25 (Bartenschlager et al. (1994) *J. Virol.* 68:5045-5055; Failla et al. (1994) *J. Virol.* 68: 3753-3760; Lin et al. (1994) 68:8147-8157; Xu et al. (1997) *J. Virol.* 71:5312-5322). The NS3 protein of both viruses also functions as a helicase (Kim et al. (1995) *Biochem. Biophys. Res. Comm.* 215: 160-166; Jin and Peterson (1995) *Arch. Biochem. Biophys.*, 323:47-53; Warrener and Collett (1995) *J. Virol.* 69:1720-1726). Finally, the NS5B

30 proteins of pestiviruses and hepaciviruses have the predicted RNA-directed RNA polymerases activity (Behrens et al.(1996) *EMBO J.* 15:12-22; Lchmannet al.(1997) *J.*

Virol. 71:8416-8428; Yuan et al.(1997) *Biochem. Biophys. Res. Comm.* 232:231-235; Hagedorn, PCT WO 97/12033; Zhong et al.(1998) *J. Virol.* 72:9365-9369).

Hepatitis C Virus

The hepatitis C virus (HCV) is the leading cause of chronic liver disease worldwide. (Boyer, N. *et al. J. Hepatol.* 32:98-112, 2000). HCV causes a slow growing viral infection and is the major cause of cirrhosis and hepatocellular carcinoma (Di Besceglie, A. M. and Bacon, B. R., *Scientific American*, Oct.: 80-85, (1999); Boyer, N. *et al. J. Hepatol.* 32:98-112, 2000). An estimated 170 million persons are infected with HCV worldwide. (Boyer, N. *et al. J. Hepatol.* 32:98-112, 2000). Cirrhosis caused by chronic hepatitis C infection accounts for 8,000-12,000 deaths per year in the United States, and HCV infection is the leading indication for liver transplantation.

HCV is known to cause at least 80% of posttransfusion hepatitis and a substantial proportion of sporadic acute hepatitis. Preliminary evidence also implicates HCV in many cases of "idiopathic" chronic hepatitis, "cryptogenic" cirrhosis, and probably hepatocellular carcinoma unrelated to other hepatitis viruses, such as Hepatitis B Virus (HBV). A small proportion of healthy persons appear to be chronic HCV carriers, varying with geography and other epidemiological factors. The numbers may substantially exceed those for HBV, though information is still preliminary; how many of these persons have subclinical chronic liver disease is unclear. (The Merck Manual, ch. 69, p. 901, 16th ed., (1992)).

HCV is an enveloped virus containing a positive-sense single-stranded RNA genome of approximately 9.4kb. The viral genome consists of a 5' untranslated region (UTR), a long open reading frame encoding a polyprotein precursor of approximately 3011 amino acids, and a short 3' UTR. The 5' UTR is the most highly conserved part of the HCV genome and is important for the initiation and control of polyprotein translation. Translation of the HCV genome is initiated by a cap-independent mechanism known as internal ribosome entry. This mechanism involves the binding of ribosomes to an RNA sequence known as the internal ribosome entry site (IRES). An RNA pseudoknot structure has recently been determined to be an essential structural

element of the HCV IRES. Viral structural proteins include a nucleocapsid core protein (C) and two envelope glycoproteins, E1 and E2.

HCV also encodes two proteinases, a zinc-dependent metalloproteinase encoded by the NS2-NS3 region and a serine proteinase encoded in the NS3 region. These proteinases are required for cleavage of specific regions of the precursor polyprotein into mature peptides: the junction between NS2 and NS3 is autocatalytically cleaved the NS2/NS3 protease, while the remaining junctions are cleaved by the N-terminal serine protease domain of NS3 complexed with NS4A. The NS3 protein contains the NTP-dependent helicase activity that unwinds duplex RNA during replication. The hydrophobic carboxy-terminal 21 amino acids of nonstructural protein 5, NS5B, contains the RNA-dependent RNA polymerase that is essential for viral replication (*Fields Virology*, Fourth Edition, Editors: Fields, B. N., Knipe, D. M., and Howley, P. M., Lippincott-Raven Publishers, Philadelphia, PA, 2001, Chapter 32, pp. 1014-1015). NS5B is known to bind RNAs nonspecifically, and to interact directly with NS3 and NS4A that, in turn, form complexes with NS4B and NS5A (*Id.* @ 1015; Ishido et al., *Biochem. Biophys. Res. Commun.*, 1998; 244:35-40). Certain *in vitro* experiments using NS5B and guanosine 5'-mono-, di-, and triphosphate as well as 5'-triphosphate of 2'-deoxy- and 2',3'-dideoxy-guanosine as HCV inhibitors suggest that HCV-RdRP may have a strict specificity for 5'-triphosphates and 2'- and 3'-OH groups (Watanabe et al., U.S. 2002/0055483). Otherwise, the function(s) of the remaining nonstructural proteins, NS4A, NS4B, and NS5A (the amino-terminal half of nonstructural protein 5) remain unknown.

A significant focus of current antiviral research is directed to the development of improved methods of treatment of chronic HCV infections in humans (Di Besceglie, A. M. and Bacon, B. R., *Scientific American*, Oct.: 80-85, (1999)).

Methods to Treat *Flaviviridae* Infections

The development of new antiviral agents for *Flaviviridae* infections, especially hepatitis C, is currently underway. Specific inhibitors of HCV-derived enzymes such as protease, helicase, and polymerase inhibitors are being developed. Drugs that inhibit other steps in HCV replication are also in development, for example, drugs that block

production of HCV antigens from the RNA (IRES inhibitors), drugs that prevent the normal processing of HCV proteins (inhibitors of glycosylation), drugs that block entry of HCV into cells (by blocking its receptor) and nonspecific cytoprotective agents that block cell injury caused by the virus infection. Further, molecular approaches are also being developed to treat hepatitis C, for example, ribozymes, which are enzymes that break down specific viral RNA molecules, and antisense oligonucleotides, which are small complementary segments of DNA that bind to viral RNA and inhibit viral replication, are under investigation. A number of HCV treatments are reviewed by Bymock *et al.* in *Antiviral Chemistry & Chemotherapy*, 11:2; 79-95 (2000) and De Francesco *et al.* in *Antiviral Research*, 58: 1-16 (2003).

Idenix Pharmaceuticals, Ltd. discloses branched nucleosides, and their use in the treatment of HCV and flaviviruses and pestiviruses in US Patent Publication Nos. 2003/0050229 A1, 2004/0097461 A1, 2004/0101535 A1, 2003/0060400 A1, 2004/0102414 A1, 2004/0097462 A1, and 2004/0063622 A1 which correspond to International Publication Nos. WO 01/90121 and WO 01/92282. A method for the treatment of hepatitis C infection (and flaviviruses and pestiviruses) in humans and other host animals is disclosed in the Idenix publications that includes administering an effective amount of a biologically active 1', 2', 3' or 4'-branched β -D or β -L nucleosides or a pharmaceutically acceptable salt or prodrug thereof, administered either alone or in combination, optionally in a pharmaceutically acceptable carrier. See also U.S. Patent Publication Nos. 2004/0006002 and 2004/0006007 as well as WO 03/026589 and WO 03/026675. Idenix Pharmaceuticals, Ltd. also discloses in US Patent Publication No. 2004/0077587 pharmaceutically acceptable branched nucleoside prodrugs, and their use in the treatment of HCV and flaviviruses and pestiviruses in prodrugs. See also PCT Publication Nos. WO 04/002422, WO 04/002999, and WO 04/003000. Further, Idenix Pharmaceuticals, Ltd. also discloses in WO 04/046331 Flaviviridae mutations caused by biologically active 2'-branched β -D or β -L nucleosides or a pharmaceutically acceptable salt or prodrug thereof.

Biota Inc. discloses various phosphate derivatives of nucleosides, including 1', 2', 3' or 4'-branched β -D or β -L nucleosides, for the treatment of hepatitis C infection in International Patent Publication WO 03/072757.

Emory University and the University of Georgia Research Foundation, Inc. (UGARF) discloses the use of 2'-fluoronucleosides for the treatment of HCV in US Patent No. 6,348,587. See also US Patent Publication No. 2002/0198171 and International Patent Publication WO 99/43691.

5 BioChem Pharma Inc. (now Shire Biochem, Inc.) discloses the use of various 1,3-dioxolane nucleosides for the treatment of a *Flaviviridae* infection in US Patent No. 6,566,365. See also US Patent Nos. 6,340,690 and 6,605,614; US Patent Publication Nos. 2002/0099072 and 2003/0225037, as well as International Publication No. WO 01/32153 and WO 00/50424..

10 BioChem Pharma Inc. (now Shire Biochem, Inc.) also discloses various other 2'-halo, 2'-hydroxy and 2'-alkoxy nucleosides for the treatment of a *Flaviviridae* infection in US Patent Publication No. 2002/0019363 as well as International Publication No. WO 01/60315 (PCT/CA01/00197; filed February 19, 2001).

15 ICN Pharmaceuticals, Inc. discloses various nucleoside analogs that are useful in modulating immune response in US Patent Nos. 6,495,677 and 6,573,248. See also WO 98/16184, WO 01/68663, and WO 02/03997.

20 US Patent No. 6,660,721; US Patent Publication Nos. 2003/083307 A1, 2003/008841 A1, and 2004/0110718; as well as International Patent Publication Nos. WO 02/18404; WO 02/100415, WO 02/094289, and WO 04/043159; filed by F. Hoffmann-La Roche AG, discloses various nucleoside analogs for the treatment of HCV RNA replication.

25 Pharmasset Limited discloses various nucleosides and antimetabolites for the treatment of a variety of viruses, including *Flaviviridae*, and in particular HCV, in US Patent Publication Nos. 2003/0087873, 2004/0067877, 2004/0082574, 2004/0067877, 2004/002479, 2003/0225029, and 2002/00555483, as well as International Patent Publication Nos. WO 02/32920, WO 01/79246, WO 02/48165, WO 03/068162, WO 03/068164 and WO 2004/013298.

30 Merck & Co., Inc. and Isis Pharmaceuticals disclose in US Patent Publication No. 2002/0147160, 2004/0072788, 2004/0067901, and 2004/0110717; as well as the corresponding International Patent Publication Nos. WO 02/057425 (PCT/US02/01531; filed January 18, 2002) and WO 02/057287 (PCT/US02/03086; filed January 18, 2002)

various nucleosides, and in particular several pyrrolopyrimidine nucleosides, for the treatment of viruses whose replication is dependent upon RNA-dependent RNA polymerase, including Flaviviridae, and in particular HCV. See also WO 2004/000858, WO 2004/003138, WO 2004/007512, and WO 2004/009020.

5 US Patent Publication No. 2003/028013 A1 as well as International Patent Publication Nos. WO 03/051899, WO 03/061576, WO 03/062255 WO 03/062256, WO 03/062257, and WO 03/061385, filed by Ribapharm, also are directed to the use of certain nucleoside analogs to treat hepatitis C virus.

10 Genelabs Technologies disclose in US Patent Publication No. 2004/0063658 as well as International Patent Publication Nos. WO 03/093290 and WO 04/028481 various base modified derivatives of nucleosides, including 1', 2', 3' or 4'-branched β -D or β -L nucleosides, for the treatment of hepatitis C infection.

15 Eldrup et al. (Oral Session V, Hepatitis C Virus, *Flaviviridae*; 16th International Conference on Antiviral Research (April 27, 2003, Savannah, Ga.) p. A75) described the structure activity relationship of 2'-modified nucleosides for inhibition of HCV.

20 Bhat et al (Oral Session V, Hepatitis C Virus, *Flaviviridae*; 16th International Conference on Antiviral Research (April 27, 2003, Savannah, Ga.); p A75) describe the synthesis and pharmacokinetic properties of nucleoside analogues as possible inhibitors of HCV RNA replication. The authors report that 2'-modified nucleosides demonstrate potent inhibitory activity in cell-based replicon assays.

Olsen et al. (Oral Session V, Hepatitis C Virus, *Flaviviridae*; 16th International Conference on Antiviral Research (April 27, 2003, Savannah, Ga.) p A76) also described the effects of the 2'-modified nucleosides on HCV RNA replication.

25 Drug-resistant variants of viruses can emerge after prolonged treatment with an antiviral agent. Drug resistance most typically occurs by mutation of a gene that encodes for an enzyme used in viral replication, and, for example, in the case of HIV, reverse transcriptase, protease, or DNA polymerase. It has been demonstrated that the efficacy of a drug against viral infection can be prolonged, augmented, or restored by
30 administering the compound in combination or alternation with a second, and perhaps

third, antiviral compound that induces a different mutation from that caused by the principle drug. Alternatively, the pharmacokinetics, biodistribution, or other parameter of the drug can be altered by such combination or alternation therapy. In general, combination therapy is typically preferred over alternation therapy because it induces multiple simultaneous pressures on the virus. One cannot predict, however, what mutations will be induced in the viral genome by a given drug, whether the mutation is permanent or transient, or how an infected cell with a mutated viral sequence will respond to therapy with other agents in combination or alternation. This is exacerbated by the fact that there is a paucity of data on the kinetics of drug resistance in long-term cell cultures treated with modern antiviral agents.

In view of the severity of diseases associated with pestiviruses, flaviviruses, and hepatitis C virus, and their pervasiveness in animals and humans, it is an object of the present invention to provide a compound, method and composition for the treatment of a host infected with any member of the family *Flaviviridae*, including hepatitis C virus.

Further, it is an object of the present invention to provide a compound, method and pharmaceutically-acceptable composition for the prophylaxis and/or treatment of a host, and particularly a human, infected with any member of the family *Flaviviridae*.

Further, given the rising threat of other *Flaviviridae* infections, there remains a strong need to provide new effective pharmaceutical agents that have low toxicity to the host.


Therefore, it is an object of the present invention to provide a compound, method and composition for the treatment of a host infected with any member of the family *Flaviviridae*, including hepatitis C virus, that have low toxicity to the host.

It is another object of the present invention to provide a compound, method and composition generally for the treatment of patients infected with *pestiviruses*, *flaviviruses*, or *hepaciviruses*.

SUMMARY OF THE INVENTION

Methods and compositions for the treatment of pestivirus, flavivirus and hepatitis C virus infection are described that include administering an effective amount of a beta-D or beta-L-nucleoside of the Formulae (I) and (II), or a pharmaceutically acceptable salt or prodrug thereof.

In a first principal embodiment, a compound of the Formula (I), or a pharmaceutically acceptable salt or prodrug thereof, is provided:



(I)

wherein

each R is independently H, phosphate (including mono-, di-, or triphosphate or a stabilized phosphate prodrug) or phosphonate; optionally substituted alkyl including lower alkyl, optionally substituted alkenyl or alkynyl, acyl, -C(O)-(alkyl), -C(O)(lower alkyl), -C(O)-(alkenyl), -C(O)-(alkynyl), lipid, phospholipid, carbohydrate, peptide, cholesterol, an amino acid residue or derivative, or other pharmaceutically acceptable leaving group that is capable of providing H or phosphate when administered *in vivo*;

n is 0-2;

when X is CH₂, CHOH, CH-alkyl, CH-alkenyl, CH-alkynyl, C-dialkyl, CH-O-alkyl, CH-O-alkenyl, CH-O-alkynyl, CH-S-alkyl, CH-S-alkenyl, CH-S-alkynyl, CH-halogen, or C-(halogen)₂,

then each R¹ and R^{1'} is independently H, OH, optionally substituted alkyl including lower alkyl, azido, cyano, optionally substituted alkenyl or alkynyl, -C(O)O-(alkyl), -C(O)O(lower alkyl), -C(O)O-(alkenyl), -C(O)O-(alkynyl), -O(acyl), -O(lower acyl),

-O(alkyl), -O(lower alkyl), -O(alkenyl), -O(alkynyl), halogen, halogenated alkyl, -NO₂, -NH₂, -NH(lower alkyl), -N(lower alkyl)₂, -NH(acyl), -N(acyl)₂, -C(O)NH₂, -C(O)NH(alkyl), -C(O)N(alkyl)₂, S(O)N-alkyl, S(O)N-alkenyl, S(O)N-alkynyl, or SCH-halogen, wherein alkyl, alkenyl, and/or alkynyl may optionally be substituted;


5 when X is O, S[O]_n, NH, N-alkyl, N-alkenyl, N-alkynyl, S(O)N-alkyl, S(O)N-alkenyl, S(O)N-alkynyl, or SCH-halogen,

then each R¹ and R^{1'} is independently H, optionally substituted alkyl including lower alkyl, azido, cyano, optionally substituted alkenyl or alkynyl, -C(O)O-(alkyl), -C(O)O(lower alkyl), -C(O)O-(alkenyl), -C(O)O-(alkynyl), halogenated alkyl,
10 -C(O)NH₂, -C(O)NH(alkyl), -C(O)N(alkyl)₂, -C(H)=N-NH₂, C(S)NH₂, C(S)NH(alkyl), or C(S)N(alkyl)₂, wherein alkyl, alkenyl, and/or alkynyl may optionally be substituted;

each R² and R³ is independently H, OH, NH₂, SH, F, Cl, Br, I, CN, NO₂, -C(O)NH₂, -C(O)NH(alkyl), and -C(O)N(alkyl)₂, N₃, optionally substituted alkyl including lower
15 alkyl, optionally substituted alkenyl or alkynyl, halogenated alkyl, -C(O)O-(alkyl), -C(O)O(lower alkyl), -C(O)O-(alkenyl), -C(O)O-(alkynyl), -O(acyl), -O(alkyl), -O(alkenyl), -O(alkynyl), -OC(O)NH₂, NC, C(O)OH, SCN, OCN, -S(alkyl), -S(alkenyl), -S(alkynyl), -NH(alkyl), -N(alkyl)₂, -NH(alkenyl), -NH(alkynyl), an amino acid residue or derivative, a prodrug or leaving group that provides OH *in vivo*, or an optionally substituted 3-7 membered heterocyclic ring having O, S and/or
20 N independently as a heteroatom taken alone or in combination;

each R^{2'} and R^{3'} is independently H; optionally substituted alkyl, alkenyl, or alkynyl; -C(O)O(alkyl), -C(O)O(lower alkyl), -C(O)O(alkenyl), -C(O)O(alkynyl), -C(O)NH₂, -C(O)NH(alkyl), -C(O)N(alkyl)₂, -O(acyl), -O(lower acyl), -O(alkyl), -O(lower
25 alkyl), -O(alkenyl), halogen, halogenated alkyl and particularly CF₃, azido, cyano, NO₂, -S(alkyl), -S(alkenyl), -S(alkynyl), NH₂, -NH(alkyl), -N(alkyl)₂, -NH(alkenyl), -NH(alkynyl), -NH(acyl), or -N(acyl)₂, and R₃ at 3'-C may also be OH; and

Base is selected from the group consisting of:



wherein:

5 each A independently is N or C-R⁵;

each W is H, OH, -O(acyl), -O(C₁₋₄ alkyl), -O(alkenyl), -O(alkynyl), -OC(O)R⁴R⁴,
 -OC(O)N R⁴R⁴, SH, -S(acyl), -S(C₁₋₄ alkyl), NH₂, NH(acyl), N(acyl)₂, NH(C₁₋₄
 alkyl), N(C₁₋₄ alkyl)₂, -N(cycloalkyl) C₁₋₄ alkylamino, di(C₁₋₄ alkyl)amino, C₃₋₆
 cycloalkylamino, halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, CN, SCN, OCN, SH, N₃, NO₂,
 10 NH=NH₂, N₃, NHOH, -C(O)NH₂, -C(O)NH(acyl), -C(O)N(acyl)₂, -C(O)NH(C₁₋₄
 alkyl), -C(O)N(C₁₋₄ alkyl)₂, -C(O)N(alkyl)(acyl), or halogenated alkyl;

each Z is O, S, NH, N-OH, N-NH₂, NH(alkyl), N(alkyl)₂, N-cycloalkyl, alkoxy, CN,
 SCN, OCN, SH, NO₂, NH₂, N₃, NH=NH, NH(alkyl), N(alkyl)₂, CONH₂,
 CONH(alkyl), or CON(alkyl)₂;

15 each R⁴ is independently H, acyl, or C₁₋₆ alkyl;


each R^5 is independently H, Cl, Br, F, I, CN, OH, optionally substituted alkyl, alkenyl or alkynyl, carboxy, $C(=NH)NH_2$, C_{1-4} alkoxy, C_{1-4} alkyloxycarbonyl, N_3 , NH_2 , $NH(alkyl)$, $N(alkyl)_2$, NO_2 , N_3 , halogenated alkyl especially CF_3 , C_{1-4} alkylamino, $di(C_{1-4} alkyl)amino$, C_{3-6} cycloalkylamino, C_{1-6} alkoxy, SH, $-S(C_{1-4} alkyl)$, $-S(C_{1-4} alkenyl)$, $-S(C_{1-4} alkynyl)$, C_{1-6} alkylthio, C_{1-6} alkylsulfonyl, $(C_{1-4} alkyl)_{0-2}$ aminomethyl, C_{3-6} cycloalkylamino -alkenyl, -alkynyl, $-(O)alkyl$, $-(O)alkenyl$, $-(O)alkynyl$, $-(O)acyl$, $-O(C_{1-4} alkyl)$, $-O(C_{1-4} alkenyl)$, $-O(C_{1-4} alkynyl)$, $-O-C(O)NH_2$, $-OC(O)N(alkyl)$, $-OC(O)R'R''$, $-C(O)OH$, $C(O)O-alkyl$, $C(O)O-alkenyl$, $C(O)O-alkynyl$, S-alkyl, S-acyl, S-alkenyl, S-alkynyl, SCN, OCN, NC, $-C(O)-NH_2$, $C(O)NH(alkyl)$, $C(O)N(alkyl)_2$, $C(O)NH(acyl)$, $C(O)N(acyl)_2$, (S)- NH_2 , NH-alkyl, $N(dialkyl)_2$, NH-acyl, N-diacyl, or a 3-7 membered heterocycle having O, S, or N taken independently in any combination;

each R' and R'' independently is H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, halogen, halogenated alkyl, OH, CN, N_3 , carboxy, C_{1-4} alkoxycarbonyl, NH_2 , C_{1-4} alkylamino, $di(C_{1-4} alkyl)amino$, C_{1-6} alkoxy, C_{1-6} alkylsulfonyl, or $(C_{1-4} alkyl)_{0-2}$ aminomethyl; and

all tautomeric, enantiomeric and stereoisomeric forms thereof;

with the caveat that when X is S in Formula (I), then the compound is not 5-(4-aminoimidazo [4,5-d][1,2,3]triazin-7-yl)-2-hydroxymethyl-tetrahydro-thiophen-3-ol or 7-(4-hydroxy-5-hydroxy-methyl-tetrahydro-thiophen-2-yl)-3,7-dihydro-imidazo[4,5-d][1,2,3]triazin-4-one.

In a second principal embodiment, a compound of the Formula (II), or a pharmaceutically acceptable salt or prodrug thereof, is provided:



(II)

wherein:

R , R^2 , $R^{2'}$, R^3 , and $R^{3'}$ are all as defined above;

X^* is CY^3 ;

5 Y^3 is hydrogen, alkyl, bromo, chloro, fluoro, iodo, azido, cyano, alkenyl, alkynyl, $-C(O)O(alkyl)$, $-C(O)O(lower\ alkyl)$, CF_3 , $-CONH_2$, $-CONH(alkyl)$, or $-CON(alkyl)_2$;

R^1 is H, OH, optionally substituted alkyl including lower alkyl, azido, cyano, optionally substituted alkenyl or alkynyl, $-C(O)O-(alkyl)$, $-C(O)O(lower\ alkyl)$, $-C(O)O-(alkenyl)$, $-C(O)O-(alkynyl)$, $-O(acyl)$, $-O(lower\ acyl)$, $-O(alkyl)$, $-O(lower\ alkyl)$, $-O(alkenyl)$, $-O(alkynyl)$, halogen, halogenated alkyl, $-NO_2$, $-NH_2$, $-NH(lower\ alkyl)$,
10 $-N(lower\ alkyl)_2$, $-NH(acyl)$, $-N(acyl)_2$, $-C(O)NH_2$, $-C(O)NH(alkyl)$, or $-C(O)N(alkyl)_2$, wherein an optional substitution on alkyl, alkenyl, and/or alkynyl may be one or more halogen, hydroxy, alkoxy or alkylthio groups taken in any combination;


Base is defined as above for formulae (A) – (G); and

15 all tautomeric, enantiomeric and stereoisomeric forms thereof;


with the caveat that when X is S in Formula (I), then the compound is not 5-(4-amino-imidazo [4,5-d][1,2,3]triazin-7-yl)-2-hydroxymethyl-tetrahydro-thiophen-3-ol or 7-(4-hydroxy-5-hydroxy-methyl-tetrahydro-thiophen-2-yl)-3,7-dihydro-imidazo[4,5-d][1,2,3]triazin-4-one.

20


In preferred embodiments, Bases (A)-(G) have a structure selected from the group consisting of:




(i)




(ii)



(iii)



(iv)



wherein

each R' and R'' independently is H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, halogen, halogenated alkyl, OH, CN, N₃, carboxy, C₁₋₄alkoxycarbonyl, NH₂, C₁₋₄ alkylamino, di(C₁₋₄ alkyl)amino, C₁₋₆ alkoxy, C₁₋₆ alkylsulfonyl, (C₁₋₄ alkyl)₀₋₂ aminomethyl, as provided above in the definitions of A and Z for the Base Formulae (A)-(G);

each W is independently H, Cl, Br, I, F, halogenated alkyl, alkoxy, OH, SH, O-alkyl, S-alkyl, O-alkenyl, O-alkynyl, S-alkenyl, S-alkynyl, -OC(O)NR⁴R⁴, O-acyl, S-acyl, CN, SCN, OCN, NO₂, N₃, NH₂, NH(alkyl), N(alkyl)₂, NH-cycloalkyl, NH-acyl, NH=NH, CONH₂, CONH(alkyl), or CON(alkyl)₂; and

each R⁴ is independently H, acyl, or C₁₋₆ alkyl;

each Z is O, S, NH, N-OH, N-NH₂, NH(alkyl), N(alkyl)₂, N-cycloalkyl, alkoxy, CN, SCN, OCN, SH, NO₂, NH₂, N₃, NH=NH, NH(alkyl), N(alkyl)₂, CONH₂, CONH(alkyl), or CON(alkyl)₂.

In its preferred embodiments, the compounds of the present invention comprise nucleosides in which each variable in Formula (I) is selected from the following, in any combination: X is O or S; R is H or phosphate; R₁ is H, CH₂OH, or CONH₂; R₂ is OH or F; R₃ is alkyl, especially methyl or propynyl, or H at the 3' position; A is H, CH or N; Z

is O, S, or NH; W is NH₂, Cl, OMe, OH, NH-cyclopropyl, S-Me; and each R' and R'' independently is Cl, CN, CONH₂ or Me.

5 In its preferred embodiments for Formula (II), the compounds of the present invention comprise nucleosides in which each variable in Formula (II) is selected from the following, in any combination: X* is CH; R is H or phosphate; R₁ is H, CH₂OH, or CONH₂; R₂ is OH or F; R₃ is alkyl, especially methyl or propynyl, or H at the 3' position; A is H, CH or N; Z is O, S, or NH; W is NH₂, Cl, OMe, OH, NH-cyclopropyl, S-Me; and each R' and R'' independently is Cl, CN, CONH₂ or Me.

10 In all embodiments, optional substituents are selected from the group consisting of one or more halogen, amino, hydroxy, carboxy and alkoxy groups or atoms, among others. It is to be understood that all stereoisomeric and tautomeric forms of the compounds shown are included herein.

15 The active compounds of the present invention can be administered in combination, alternation or sequential steps with another anti-HCV agent. In combination therapy, effective dosages of two or more agents are administered together, whereas in alternation or sequential-step therapy, an effective dosage of each agent is administered serially or sequentially. The dosages given will depend on absorption, inactivation and excretion rates of the drug as well as other factors known to those of skill in the art. It is to be noted that dosage values will also vary with the severity of the condition to be alleviated. It is to be further understood that for any particular subject, specific dosage regimens and schedules should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions.

20

BRIEF DESCRIPTION OF THE DRAWINGS

25 **Figure 1** show generalized structural depictions for Formula (I) and Formula (II) of the ribofuranosynucleosides of the present invention.

Figure 2 shows generalized structures for the 2-azapurine bases of the present invention.

Figure 3 shows structural depictions for preferred bases of the present invention.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides a compound, method and composition for the treatment of a pestivirus, flavivirus and/or hepatitis C in humans or other host animals that includes administering an effective anti-pestivirus, anti-flavivirus or anti-HCV treatment amount of a beta-D- or beta-L-nucleoside as described herein, or a pharmaceutically acceptable salt or prodrug thereof, optionally in a pharmaceutically acceptable carrier. The compounds of this invention either possess antiviral activity, or are metabolized to a compound that exhibits such activity.


Flaviviruses included within the scope of this invention are discussed generally in *Fields Virology*, Editors: Fields, N., Knipe, D.M. and Howley, P.M.; Lippincott-Raven Publishers, Philadelphia, PA; Chapter 31 (1996). Specific flaviviruses include, without limitation: Absettarov; Alfuy; Apoi; Aroa; Bagaza; Banzi; Bououi; Bussuquara; Cacipacore; Carey Island; Dakar bat; Dengue viruses 1, 2, 3 and 4; Edge Hill; Entebbe bat; Gadgets Gully; Hanzalova; Hypr; Ilheus; Israel turkey meningoencephalitis; Japanese encephalitis; Jugra; Jutiapa; Kadam; Karshi; Kedougou; Kokoera; Koutango; Kumlinge; Kunjin; Kyasanur Forest disease; Langat; Louping ill; Meaban; Modoc; Montana myotis leukoencephalitis; Murray valley encephalitis; Naranjal; Negishi; Ntaya; Omsk hemorrhagic fever; Phnom-Penh bat; Powassan; Rio Bravo; Rocio; Royal Farm; Russian spring-summer encephalitis; Saboya; St. Louis encephalitis; Sal Vieja; San Perlita; Saumarez Reef; Sepik; Sokuluk; Spondweni; Stratford; Temusu; Tyuleny; Uganda S, Usutu, Wesselsbron; West Nile; Yaounde; Yellow fever; and Zika.

Pestiviruses included within the scope of this invention are also discussed generally in *Fields Virology* (Id.). Specific pestiviruses include, without limitation: bovine viral diarrhea virus ("VDV"); classical swine fever virus ("CSFV") also known as hog cholera virus; and border disease virus ("DV").

HCV is a member of the family, *Flaviviridae*; however, HCV now has been placed in a new monotypic genus, *hepacivirus*.

Active Compounds, Pharmaceutically Acceptable Salts and Prodrugs Thereof

In a first principal embodiment, a compound of the Formula (I), or a pharmaceutically acceptable salt or prodrug thereof, is provided:



(I)

wherein

R is H, phosphate (including mono-, di-, or triphosphate or a stabilized phosphate prodrug) or phosphonate; optionally substituted alkyl including lower alkyl, optionally substituted alkenyl or alkynyl, acyl, -C(O)-(alkyl), -C(O)(lower alkyl), -C(O)-(alkenyl), -C(O)-(alkynyl), lipid, phospholipid, carbohydrate, peptide, cholesterol, an amino acid residue or derivative, or other pharmaceutically acceptable leaving group that is capable of providing H or phosphate when administered *in vivo*;

n is 0-2;

when X is CH₂, CHOH, CH-alkyl, CH-alkenyl, CH-alkynyl, C-dialkyl, CH-O-alkyl, CH-O-alkenyl, CH-O-alkynyl, CH-S-alkyl, CH-S-alkenyl, CH-S-alkynyl, CH-halogen, or C-(halogen)₂,

then each R¹ and R^{1'} is independently H, OH, optionally substituted alkyl including lower alkyl, azido, cyano, optionally substituted alkenyl or alkynyl, -C(O)O-(alkyl), -C(O)O(lower alkyl), -C(O)O-(alkenyl), -C(O)O-(alkynyl), -O(acyl), -O(lower acyl), -O(alkyl), -O(lower alkyl), -O(alkenyl), -O(alkynyl), halogen, halogenated alkyl, -NO₂, -NH₂, -NH(lower alkyl), -N(lower alkyl)₂, -NH(acyl), -N(acyl)₂, -C(O)NH₂, -C(O)NH(alkyl), -C(O)N(alkyl)₂, S(O)N-alkyl, S(O)N-alkenyl, S(O)N-alkynyl, or SCH-halogen, wherein alkyl, alkenyl, and/or alkynyl may optionally be substituted;


when X is O, S[O]_n, NH, N-alkyl, N-alkenyl, N-alkynyl, S(O)N-alkyl, S(O)N-alkenyl, S(O)N-alkynyl, or SCH-halogen,


then each R^1 and $R^{1'}$ is independently H, optionally substituted alkyl including lower alkyl, azido, cyano, optionally substituted alkenyl or alkynyl, $-C(O)O-(alkyl)$, $-C(O)O(lower\ alkyl)$, $-C(O)O-(alkenyl)$, $-C(O)O-(alkynyl)$, halogenated alkyl, $-C(O)NH_2$, $-C(O)NH(alkyl)$, $-C(O)N(alkyl)_2$, $-C(H)=N-NH_2$, $C(S)NH_2$, $C(S)NH(alkyl)$, or $C(S)N(alkyl)_2$, wherein alkyl, alkenyl, and/or alkynyl may optionally be substituted;

each R^2 and R^3 is independently is OH, NH_2 , SH, F, Cl, Br, I, CN, NO_2 , $-C(O)NH_2$, $-C(O)NH(alkyl)$, $-C(O)N(alkyl)_2$, N_3 , optionally substituted alkyl including lower alkyl, optionally substituted alkenyl or alkynyl, halogenated alkyl, $-C(O)O-(alkyl)$, $-C(O)O(lower\ alkyl)$, $-C(O)O-(alkenyl)$, $-C(O)O-(alkynyl)$, $-O(acyl)$, $-O(alkyl)$, $-O(alkenyl)$, $-O(alkynyl)$, $-OC(O)NH_2$, NC, $C(O)OH$, SCN, OCN, $-S(alkyl)$, $-S(alkenyl)$, $-S(alkynyl)$, $-NH(alkyl)$, $-N(alkyl)_2$, $-NH(alkenyl)$, $-NH(alkynyl)$, an amino acid residue or derivative, a prodrug or leaving group that provides OH *in vivo*, or an optionally substituted 3-7 membered heterocyclic ring having O, S and/or N independently as a heteroatom taken alone or in combination;

each $R^{2'}$ and $R^{3'}$ independently is H; optionally substituted alkyl, alkenyl, or alkynyl; $-C(O)O(alkyl)$, $-C(O)O(lower\ alkyl)$, $-C(O)O(alkenyl)$, $-C(O)O(alkynyl)$, $-C(O)NH_2$, $-C(O)NH(alkyl)$, $-C(O)N(alkyl)_2$, $-O(acyl)$, $-O(lower\ acyl)$, $-O(alkyl)$, $-O(lower\ alkyl)$, $-O(alkenyl)$, halogen, halogenated alkyl and particularly CF_3 , azido, cyano, NO_2 , $-S(alkyl)$, $-S(alkenyl)$, $-S(alkynyl)$, NH_2 , $-NH(alkyl)$, $-N(alkyl)_2$, $-NH(alkenyl)$, $-NH(alkynyl)$, $-NH(acyl)$, or $-N(acyl)_2$, and R_3 at 3'-C may also be OH; and

Base is selected from the group consisting of:





wherein

each A independently is N or C-R⁵;

- 5 W is H, OH, -O(acyl), -O(C₁₋₄ alkyl), -O(alkenyl), -O(alkynyl), -OC(O)R⁴R⁴, -OC(O)N R⁴R⁴, SH, -S(acyl), -S(C₁₋₄ alkyl), NH₂, NH(acyl), N(acyl)₂, NH(C₁₋₄ alkyl), N(C₁₋₄ alkyl)₂, -N(cycloalkyl) C₁₋₄ alkylamino, di(C₁₋₄ alkyl)amino, C₃₋₆ cycloalkylamino, halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, CN, SCN, OCN, SH, N₃, NO₂, NH=NH₂, N₃, NHOH, -C(O)NH₂, -C(O)NH(acyl), -C(O)N(acyl)₂, -C(O)NH(C₁₋₄ alkyl),
 10 -C(O)N(C₁₋₄ alkyl)₂, -C(O)N(alkyl)(acyl), or halogenated alkyl;

Z is O, S, NH, N-OH, N-NH₂, NH(alkyl), N(alkyl)₂, N-cycloalkyl, alkoxy, CN, SCN, OCN, SH, NO₂, NH₂, N₃, NH=NH, NH(alkyl), N(alkyl)₂, CONH₂, CONH(alkyl), or CON(alkyl)₂;

each R⁴ is independently H, acyl, or C₁₋₆ alkyl;

- 15 each R⁵ is independently H, Cl, Br, F, I, CN, OH, optionally substituted alkyl, alkenyl or alkynyl, carboxy, C(=NH)NH₂, C₁₋₄ alkoxy, C₁₋₄ alkyloxycarbonyl, N₃, NH₂, NH(alkyl), N(alkyl)₂, NO₂, N₃, halogenated alkyl especially CF₃, C₁₋₄ alkylamino, di(C₁₋₄ alkyl)amino, C₃₋₆ cycloalkylamino, C₁₋₆ alkoxy, SH, -S(C₁₋₄ alkyl), -S(C₁₋₄ alkenyl), -S(C₁₋₄ alkynyl), C₁₋₆ alkylthio, C₁₋₆ alkylsulfonyl, (C₁₋₄ alkyl)₀₋₂ aminomethyl, C₃₋₆ cycloalkylamino -alkenyl, -alkynyl, -(O)alkyl, -(O)alkenyl,
 20

5
 10
 15
 20
 25
 30
 35
 40
 45
 50
 55
 60
 65
 70
 75
 80
 85
 90
 95
 100
 105
 110
 115
 120
 125
 130
 135
 140
 145
 150
 155
 160
 165
 170
 175
 180
 185
 190
 195
 200
 205
 210
 215
 220
 225
 230
 235
 240
 245
 250
 255
 260
 265
 270
 275
 280
 285
 290
 295
 300
 305
 310
 315
 320
 325
 330
 335
 340
 345
 350
 355
 360
 365
 370
 375
 380
 385
 390
 395
 400
 405
 410
 415
 420
 425
 430
 435
 440
 445
 450
 455
 460
 465
 470
 475
 480
 485
 490
 495
 500
 505
 510
 515
 520
 525
 530
 535
 540
 545
 550
 555
 560
 565
 570
 575
 580
 585
 590
 595
 600
 605
 610
 615
 620
 625
 630
 635
 640
 645
 650
 655
 660
 665
 670
 675
 680
 685
 690
 695
 700
 705
 710
 715
 720
 725
 730
 735
 740
 745
 750
 755
 760
 765
 770
 775
 780
 785
 790
 795
 800
 805
 810
 815
 820
 825
 830
 835
 840
 845
 850
 855
 860
 865
 870
 875
 880
 885
 890
 895
 900
 905
 910
 915
 920
 925
 930
 935
 940
 945
 950
 955
 960
 965
 970
 975
 980
 985
 990
 995


-(O)alkynyl, -(O)acyl, -O(C₁₋₄ alkyl), -O(C₁₋₄ alkenyl), -O(C₁₋₄ alkynyl), -O-C(O)NH₂, -OC(O)N(alkyl), -OC(O)R'R'', -C(O)OH, C(O)O-alkyl, C(O)O-alkenyl, C(O)O-alkynyl, S-alkyl, S-acyl, S-alkenyl, S-alkynyl, SCN, OCN, NC, -C(O)-NH₂, C(O)NH(alkyl), C(O)N(alkyl)₂, C(O)NH(acyl), C(O)N(acyl)₂, (S)-NH₂, NH-alkyl, N(dialkyl)₂, NH-acyl, N-diacyl, or a 3-7 membered heterocycle having O, S, or N taken independently in any combination;

each R' and R'' independently is H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, halogen, halogenated alkyl, OH, CN, N₃, carboxy, C₁₋₄alkoxycarbonyl, NH₂, C₁₋₄ alkylamino, di(C₁₋₄ alkyl)amino, C₁₋₆ alkoxy, C₁₋₆ alkylsulfonyl, or (C₁₋₄ alkyl)₀₋₂ aminomethyl;

all tautomeric, enantiomeric and stereoisomeric forms thereof;

with the caveat that when X is S in Formula (I), then the compound is not 5-(4-aminoimidazo [4,5-d][1,2,3]triazin-7-yl)-2-hydroxymethyl-tetrahydro-thiophen-3-ol or 7-(4-hydroxy-5-hydroxy-methyl-tetrahydro-thiophen-2-yl)-3,7-dihydro-imidazo[4,5-d][1,2,3]triazin-4-one.

In a second principal embodiment, a compound of the Formula (II), or a pharmaceutically acceptable salt or prodrug thereof, is provided:



(II)

wherein:

R, R², R^{2'}, R³, and R^{3'} are all as defined above;

X* is CY³;

Y^3 is hydrogen, alkyl, bromo, chloro, fluoro, iodo, azido, cyano, alkenyl, alkynyl, -C(O)O(alkyl), -C(O)O(lower alkyl), CF_3 , -CONH₂, -CONH(alkyl), or -CON(alkyl)₂;

R^1 is H, OH, optionally substituted alkyl including lower alkyl, azido, cyano, optionally substituted alkenyl or alkynyl, -C(O)O-(alkyl), -C(O)O(lower alkyl), -C(O)O-(alkenyl), -C(O)O-(alkynyl), -O(acyl), -O(lower acyl), -O(alkyl), -O(lower alkyl), -O(alkenyl), -O(alkynyl), halogen, halogenated alkyl, -NO₂, -NH₂, -NH(lower alkyl), -N(lower alkyl)₂, -NH(acyl), -N(acyl)₂, -C(O)NH₂, -C(O)NH(alkyl), or -C(O)N(alkyl)₂, wherein an optional substitution on alkyl, alkenyl, and/or alkynyl may be one or more halogen, hydroxy, alkoxy or alkylthio groups taken in any combination;





Base is defined as above for formulae (A) – (G); and


A and Z are as defined above,

with the caveat that when X is S in Formula (I), then the compound is not 5-(4-aminoimidazo [4,5-d][1,2,3]triazin-7-yl)-2-hydroxymethyl-tetrahydro-thiophen-3-ol or 7-(4-hydroxy-5-hydroxy-methyl-tetrahydro-thiophen-2-yl)-3,7-dihydro-imidazo[4,5-d][1,2,3]triazin-4-one; and

all tautomeric, enantiomeric and stereoisomeric forms thereof.

In preferred embodiments, Bases (A)-(G) have a structure selected from the group consisting of:





wherein

each R' and R'' independently is H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, halogen, halogenated alkyl, OH, CN, N₃, carboxy, C₁₋₄alkoxycarbonyl, NH₂, C₁₋₄ alkylamino, di(C₁₋₄ alkyl)amino, C₁₋₆ alkoxy, C₁₋₆ alkylsulfonyl, (C₁₋₄ alkyl)₀₋₂ aminomethyl, as provided above in the definitions of A and Z for the Base Formulae (A)-(G);

each W is Cl, Br, I, F, halogenated alkyl, alkoxy, OH, SH, O-alkyl, S-alkyl, O-alkenyl, O-alkynyl, S-alkenyl, S-alkynyl, -OC(O)NR⁴R⁴, O-acyl, S-acyl, CN, SCN, OCN, NO₂, N₃, NH₂, NH(alkyl), N(alkyl)₂, NH-cycloalkyl, NH-acyl, NH=NH, CONH₂, CONH(alkyl), or CON(alkyl)₂;

each R⁴ is independently H, acyl, or C₁₋₆ alkyl; and

each Z is O, S, NH, N-OH, N-NH₂, NH(alkyl), N(alkyl)₂, N-cycloalkyl, alkoxy, CN, SCN, OCN, SH, NO₂, NH₂, N₃, NH=NH, NH(alkyl), N(alkyl)₂, CONH₂, CONH(alkyl), or CON(alkyl)₂.


In its preferred embodiments, the compounds of the present invention comprise nucleosides in which each variable in Formula (I) is selected from the following, in any combination: X is O or S; R is H or phosphate; R₁ is H, CH₂OH, or CONH₂; R₂ is OH or F; R₃ is alkyl, especially methyl or propynyl, or H at the 3' position; A is H, CH or N; Z

is O, S, or NH; W is NH₂, Cl, OMe, OH, NH-cyclopropyl, S-Me; and each R' and R'' independently is Cl, CN, CONH₂ or Me.

5 In its preferred embodiments for Formula (II), the compounds of the present invention comprise nucleosides in which each variable in Formula (II) is selected from the following, in any combination: X* is CH; R is H or phosphate; R₁ is H, CH₂OH, or CONH₂; R₂ is OH or F; R₃ is alkyl, especially methyl or propynyl, or H at the 3' position; A is H, CH or N; Z is O, S, or NH; W is NH₂, Cl, OMe, OH, NH-cyclopropyl, S-Me; and each R' and R'' independently is Cl, CN, CONH₂ or Me.

10 In all embodiments, optional substituents are selected from the group consisting of one or more halogen, amino, hydroxy, carboxy and alkoxy groups or atoms, among others. It is to be understood that all stereoisomeric and tautomeric forms of the compounds shown are included herein.

In one particular embodiment, a compound of the Formula (III), or a pharmaceutically acceptable salt or prodrug thereof, is provided:



20 each R, R^{2*}, and R^{3*} independently is H, phosphate (including mono-, di-, or triphosphate or a stabilized phosphate prodrug) or phosphonate; optionally substituted alkyl including lower alkyl, optionally substituted alkenyl or alkynyl, acyl, -C(O)-(alkyl), -C(O)(lower alkyl), -C(O)-(alkenyl), -C(O)-(alkynyl), lipid, phospholipid, carbohydrate, peptide, cholesterol, an amino acid residue or derivative, or other pharmaceutically acceptable leaving group that is capable of providing H or phosphate when administered *in vivo*;

25 X is O, S[O]_n, CH₂, CHOH, CH-alkyl, CH-alkenyl, CH-alkynyl, C-dialkyl, CH-O-alkyl, CH-O-alkenyl, CH-O-alkynyl, CH-S-alkyl, CH-S-alkenyl, CH-S-alkynyl, NH, N-alkyl, N-alkenyl, N-alkynyl, S(O)N-alkyl, S(O)N-alkenyl, S(O)N-alkynyl, SCH-

halogen, or C-(halogen)₂, wherein alkyl, alkenyl or alkynyl optionally may be substituted;

n is 0-2;

5 each R^{2'} independently is H; optionally substituted alkyl, alkenyl, or alkynyl; -C(O)O(alkyl), -C(O)O(lower alkyl), -C(O)O(alkenyl), -C(O)O(alkynyl), -C(O)NH₂, -C(O)NH(alkyl), -C(O)N(alkyl)₂, -OH, -O(acyl), -O(lower acyl), -O(alkyl), -O(lower alkyl), -O(alkenyl), halogen, halogenated alkyl and particularly CF₃, azido, cyano, NO₂, -S(alkyl), -S(alkenyl), -S(alkynyl), NH₂, -NH(alkyl), -N(alkyl)₂, -NH(alkenyl), -NH(alkynyl), -NH(acyl), or -N(acyl)₂; and


10 Base is defined as above for formulae (A) – (G); and preferably is a Base as defined by structures (i) – (xi) above.

15 In one embodiment, the R^{2'} is an optionally substituted alkyl, alkenyl, or alkynyl; halogen, halogenated alkyl and particularly CF₃, azido, or cyano. In a particular embodiment, R^{2'} is an optionally substituted alkyl, alkenyl, or alkynyl; halogen, halogenated alkyl and particularly CF₃. In yet another particular embodiment, R^{2'} is CH₃ or CF₃.

20 In one embodiment, each R, R^{2*}, and R^{3*} is independently H, phosphate (including mono-, di-, or triphosphate or a stabilized phosphate prodrug) or phosphonate. In another embodiment, each R, R^{2*}, and R^{3*} is independently H. In yet another embodiment, each R, R^{2*}, and R^{3*} is independently H, acyl, or an amino acid acyl residue.

In one embodiment, X is O or S. In another embodiment, X is O.

In another particular embodiment, a compound of the Formula (IV), or a pharmaceutically acceptable salt or prodrug thereof, is provided:



(IV)

5 each R, R^{2*}, and R^{3*} independently is H, phosphate (including mono-, di-, or triphosphate or a stabilized phosphate prodrug) or phosphonate; optionally substituted alkyl including lower alkyl, optionally substituted alkenyl or alkynyl, acyl, -C(O)-(alkyl), -C(O)(lower alkyl), -C(O)-(alkenyl), -C(O)-(alkynyl), lipid, phospholipid, carbohydrate, peptide, cholesterol, an amino acid residue or derivative,
 10 or other pharmaceutically acceptable leaving group that is capable of providing H or phosphate when administered *in vivo*;

X is O, S[O]_n, CH₂, CHOH, CH-alkyl, CH-alkenyl, CH-alkynyl, C-dialkyl, CH-O-alkyl, CH-O-alkenyl, CH-O-alkynyl, CH-S-alkyl, CH-S-alkenyl, CH-S-alkynyl, NH, N-alkyl, N-alkenyl, N-alkynyl, S(O)N-alkyl, S(O)N-alkenyl, S(O)N-alkynyl, SCH-halogen, or C-(halogen)₂, wherein alkyl, alkenyl or alkynyl optionally may be
 15 substituted;

n is 0-2;

each R^{3'} independently is H; optionally substituted alkyl, alkenyl, or alkynyl; -C(O)O(alkyl), -C(O)O(lower alkyl), -C(O)O(alkenyl), -C(O)O(alkynyl), -C(O)NH₂,
 20 -C(O)NH(alkyl), -C(O)N(alkyl)₂, -OH, -O(acyl), -O(lower acyl), -O(alkyl), -O(lower alkyl), -O(alkenyl), halogen, halogenated alkyl and particularly CF₃, azido, cyano, NO₂, -S(alkyl), -S(alkenyl), -S(alkynyl), NH₂, -NH(alkyl), -N(alkyl)₂, -NH(alkenyl), -NH(alkynyl), -NH(acyl), or -N(acyl)₂; and

Base is defined as above for formulae (A) – (G); and preferably is a Base as defined by
 25 structures (i) – (xi) above.

In one embodiment, the R^{3'} is an optionally substituted alkyl, alkenyl, or alkynyl; halogen, halogenated alkyl and particularly CF₃, azido, or cyano. In a particular embodiment, R^{3'} is an optionally substituted alkyl, alkenyl, or alkynyl; halogen, halogenated alkyl and particularly CF₃. In yet another particular embodiment, R^{3'} is CH₃ or CF₃.

In one embodiment, each R, R^{2*}, and R^{3*} is independently H, phosphate (including mono-, di-, or triphosphate or a stabilized phosphate prodrug) or phosphonate. In another embodiment, each R, R^{2*}, and R^{3*} is independently H. In yet another embodiment, each R, R^{2*}, and R^{3*} is independently H, acyl, or an amino acid acyl residue.

In one embodiment, X is O or S. In another embodiment, X is O.

The beta-D- and beta-L-nucleosides of this invention belong to a class of anti-pestivirus, anti-flavivirus and anti-HCV agents that inhibit viral polymerase. Triphosphate nucleosides can be screened for their ability to inhibit viral polymerase, whether HCV, flavivirus or pestivirus, *in vitro* according to screening methods set forth below. Chiron Corporation developed a replicon system for testing potential anti-HCV compounds that utilizes a particular peptide sequence having an HCV protease-recognition site (U.S. 6,436,666; U.S. 6,416,946; U.S. 6,416,944; U.S. 6,379,886; and U.S. 6,326,151, to Chiron Corporation). Other systems for assessing the ability of compounds to inhibit HCV and related viruses include those of Rice (see U.S. 5,874,565) and the polymerase inhibition assay of Dr. Ralf Bartenschlager (see EP 1 043 399 A2).

An alternative means of assessing a compound's ability to inhibit HCV, pestivirus and/or flavivirus is through the use of predictive animal model systems. The model of choice for testing HCV is the chimpanzee, which has been used by the applicants. Chimpanzees provide an excellent mammalian system for study of anti-HCV compounds and an insight into the predictability or unpredictability of drug activity based on the closeness of their species relationship to humans.

The active compounds of the present invention can be administered in combination, alternation or sequential steps with another anti-HCV agent. In combination therapy, effective dosages of two or more agents are administered together,

whereas in alternation or sequential-step therapy, an effective dosage of each agent is administered serially or sequentially. The dosages given will depend on absorption, inactivation and excretion rates of the drug as well as other factors known to those of skill in the art. It is to be noted that dosage values will also vary with the severity of the condition to be alleviated. It is to be further understood that for any particular subject, specific dosage regimens and schedules should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions.

In particular, the present invention provides the following:

- a) a beta-D- or beta-L-nucleoside compound of Formula (I) – (IV), or a pharmaceutically acceptable salt or prodrug thereof;
- b) a pharmaceutical composition comprising a beta-D- or beta-L-nucleoside compound of Formula (I) – (IV), or a pharmaceutically acceptable salt or prodrug thereof, optionally together with a pharmaceutically acceptable carrier, excipient or diluent;
- c) a pharmaceutical composition comprising a beta-D- or beta-L-nucleoside compound of Formula (I) – (IV), or a pharmaceutically acceptable salt or prodrug thereof, with one or more other effective antiviral agents, optionally with a pharmaceutically acceptable carrier or diluent;
- d) a pharmaceutical composition for the treatment or prophylaxis of a pestivirus, flavivirus or HCV infection in a host, especially a host diagnosed as having or being at risk for such infection, comprising a beta-D- or beta-L-nucleoside compound of Formula (I) – (IV), or a pharmaceutically acceptable salt or prodrug thereof, together with a pharmaceutically acceptable carrier or diluent;
- e) a pharmaceutical formulation comprising the beta-D- or beta-L-nucleoside compound of Formula (I) – (IV), or a pharmaceutically acceptable salt or prodrug thereof, together with a pharmaceutically acceptable carrier, excipient or diluent;
- f) a method for the treatment of a pestivirus, flavivirus or HCV infection in a host comprising a beta-D- or beta-L-nucleoside compound of Formula (I) – (IV), or a pharmaceutically acceptable salt or prodrug thereof, optionally with a pharmaceutically acceptable carrier, excipient or diluent;

- g) a method for the treatment of a pestivirus, flavivirus or HCV infection in a host comprising administering an effective amount of a beta-D- or beta-L-nucleoside compound of Formula (I) – (IV), or a pharmaceutically acceptable salt or prodrug thereof, with one or more other effective antiviral agents, optionally with a pharmaceutically acceptable carrier, excipient or diluent;
- h) a method for the treatment of a pestivirus, flavivirus or HCV infection in a host comprising administering an effective amount of a beta-D- or beta-L-nucleoside compound of Formula (I) – (IV), or a pharmaceutically acceptable salt or prodrug thereof, with one or more other effective antiviral agents, optionally with a pharmaceutically acceptable carrier, excipient or diluent;
- i) a method for the treatment of a pestivirus, flavivirus or HCV infection in a host comprising administering an effective amount of a beta-D- or beta-L-nucleoside compound of Formula (I) – (IV), or a pharmaceutically acceptable salt or prodrug thereof, with one or more other effective antiviral agents, optionally with a pharmaceutically acceptable carrier, excipient or diluent;
- j) a method for the treatment of a pestivirus, flavivirus or HCV infection in a host comprising administering an effective amount of a beta-D- or beta-L-nucleoside compound of Formula (I) – (IV), or a pharmaceutically acceptable salt or prodrug thereof, with one or more other effective antiviral agents, optionally with a pharmaceutically acceptable carrier, excipient or diluent;
- k) use of a beta-D- or beta-L-nucleoside compound of Formula (I) – (IV), or a pharmaceutically acceptable salt or prodrug thereof, optionally with a pharmaceutically acceptable carrier or diluent, for the treatment of a pestivirus, flavivirus or HCV infection in a host;
- l) use of a beta-D- or beta-L-nucleoside compound of Formula (I) – (IV), or a pharmaceutically acceptable salt or prodrug thereof, with one or more other effective antiviral agents, optionally with a pharmaceutically acceptable carrier or diluent, for the treatment of a pestivirus, flavivirus and/or HCV infection in a host;
- m) use of a beta-D- or beta-L-nucleoside compound of Formula (I) – (IV), or a pharmaceutically acceptable salt or prodrug thereof, optionally with a

pharmaceutically acceptable carrier or diluent, in the manufacture of a medicament for the treatment of a pestivirus, flavivirus and/or HCV infection in a host;

- 5 n) use of a beta-D- or beta-L-nucleoside compound of Formula (I) – (IV), or a pharmaceutically acceptable salt or prodrug thereof, with one or more other effective antiviral agents and optionally with a pharmaceutically acceptable carrier, excipient or diluent, in the manufacture of a medicament for the treatment of a pestivirus, flavivirus and/or HCV infection in a host;
- 10 o) a beta-D- or beta-L-nucleoside compound of Formula (I) – (IV), or a pharmaceutically acceptable salt or prodrug thereof, substantially in the absence of enantiomers of the described nucleoside, or substantially isolated from other chemical entities;
- p) a process for the preparation of a beta-D- or beta-L-nucleoside compound of Formula (I) – (IV), or a pharmaceutically acceptable salt or prodrug thereof, as provided in more detail below; and
- 15 q) a process for the preparation of a beta-D- or beta-L-nucleoside compound of Formula (I) – (IV), or a pharmaceutically acceptable salt or prodrug thereof, substantially in the absence of enantiomers of the described nucleoside or substantially isolated from other chemical entities.

20 The active compound can be administered as any salt or prodrug that upon administration to the recipient is capable of providing directly or indirectly the parent compound, or that exhibits activity itself. Non-limiting examples are the pharmaceutically acceptable salts, which are alternatively referred to as “physiologically acceptable salts”, and a compound that has been alkylated or acylated at the 5'-position

25 or on the purine or pyrimidine base, thereby forming a type of “pharmaceutically acceptable prodrug”. Further, the modifications can affect the biological activity of the compound, in some cases increasing the activity over the parent compound. This can easily be assessed by preparing the salt or prodrug and testing its antiviral activity according to the methods described herein, or other methods known to those skilled in

30 the art.

Stereochemistry

It is appreciated that nucleosides of the present invention have several chiral centers and may exist in and be isolated in optically active and racemic forms. Some compounds may exhibit polymorphism. It is to be understood that the present invention encompasses any racemic, optically-active, diastereomeric, polymorphic, or stereoisomeric form, or mixtures thereof, of a compound of the invention, which possess the useful properties described herein. It being well known in the art how to prepare optically active forms (for example, by resolution of the racemic form by recrystallization techniques, by synthesis from optically-active starting materials, by chiral synthesis, or by chromatographic separation using a chiral stationary phase).

Examples of methods to obtain optically active materials are known in the art, and include at least the following.

- i) physical separation of crystals - a technique whereby macroscopic crystals of the individual enantiomers are manually separated. This technique can be used if crystals of the separate enantiomers exist, i.e., the material is a conglomerate, and the crystals are visually distinct;
- ii) simultaneous crystallization - a technique whereby the individual enantiomers are separately crystallized from a solution of the racemate, possible only if the latter is a conglomerate in the solid state;
- iii) enzymatic resolutions - a technique whereby partial or complete separation of a racemate by virtue of differing rates of reaction for the enantiomers with an enzyme;
- iv) enzymatic asymmetric synthesis - a synthetic technique whereby at least one step of the synthesis uses an enzymatic reaction to obtain an enantiomerically pure or enriched synthetic precursor of the desired enantiomer;
- v) chemical asymmetric synthesis - a synthetic technique whereby the desired enantiomer is synthesized from an achiral precursor

under conditions that produce asymmetry (i.e., chirality) in the product, which may be achieved using chiral catalysts or chiral auxiliaries;

- 5
- vi) diastereomer separations - a technique whereby a racemic compound is reacted with an enantiomerically pure reagent (the chiral auxiliary) that converts the individual enantiomers to diastereomers. The resulting diastereomers are then separated by chromatography or crystallization by virtue of their now more distinct structural differences and the chiral auxiliary later removed to obtain the desired enantiomer;
- 10
- vii) first- and second-order asymmetric transformations - a technique whereby diastereomers from the racemate equilibrate to yield a preponderance in solution of the diastereomer from the desired enantiomer or where preferential crystallization of the diastereomer from the desired enantiomer perturbs the equilibrium such that eventually in principle all the material is converted to the crystalline diastereomer from the desired enantiomer. The desired enantiomer is then released from the diastereomer;
- 15
- viii) kinetic resolutions - this technique refers to the achievement of partial or complete resolution of a racemate (or of a further resolution of a partially resolved compound) by virtue of unequal reaction rates of the enantiomers with a chiral, non-racemic reagent or catalyst under kinetic conditions;
- 20
- ix) enantiospecific synthesis from non-racemic precursors - a synthetic technique whereby the desired enantiomer is obtained from non-chiral starting materials and where the stereochemical integrity is not or is only minimally compromised over the course of the synthesis;
- 25
- x) chiral liquid chromatography - a technique whereby the enantiomers of a racemate are separated in a liquid mobile phase by virtue of their differing interactions with a stationary phase.
- 30

The stationary phase can be made of chiral material or the mobile phase can contain an additional chiral material to provoke the differing interactions;

- 5 xi) chiral gas chromatography - a technique whereby the racemate is volatilized and enantiomers are separated by virtue of their differing interactions in the gaseous mobile phase with a column containing a fixed non-racemic chiral adsorbent phase;
- 10 xii) extraction with chiral solvents - a technique whereby the enantiomers are separated by virtue of preferential dissolution of one enantiomer into a particular chiral solvent;
- 15 xiii) transport across chiral membranes - a technique whereby a racemate is placed in contact with a thin membrane barrier. The barrier typically separates two miscible fluids, one containing the racemate, and a driving force such as concentration or pressure differential causes preferential transport across the membrane barrier. Separation occurs as a result of the non-racemic chiral nature of the membrane which allows only one enantiomer of the racemate to pass through.

Definitions

- 20 The term "alkyl" as used herein, unless otherwise specified, refers to a saturated straight, branched, or cyclic, primary, secondary, or tertiary hydrocarbon of typically C₁ to C₁₀, and specifically includes methyl, trifluoromethyl, ethyl, propyl, isopropyl, cyclopropyl, butyl, isobutyl, *t*-butyl, pentyl, cyclopentyl, isopentyl, neopentyl, hexyl, isohexyl, cyclohexyl, cyclohexylmethyl, 3-methylpentyl, 2,2-dimethylbutyl, and 2,3-
- 25 dimethylbutyl. The term includes both substituted and unsubstituted alkyl groups. Moieties with which the alkyl group can be substituted with one or more substituents are selected from the group consisting of halo, including Cl, F, Br and I so as to form, for eg., CF₃, 2-Br-ethyl, CH₂F, CH₂Cl, CH₂CF₃, or CF₂CF₃; hydroxyl, for eg. CH₂OH; amino, for eg., CH₂NH₂, CH₂NHCH₃, or CH₂N(CH₃)₂; carboxylate; carboxamido;
- 30 alkylamino; arylamino; alkoxy; aryloxy; nitro; azido, for eg., CH₂N₃; cyano, for eg.,

CH₂CN; thio; sulfonic acid; sulfate; phosphonic acid; phosphate; and phosphonate, either unprotected or protected as necessary, known to those skilled in the art, for eg., as taught in Greene et al., *Protective Groups in Organic Synthesis*, John Wiley and Sons, Second Edition (1991), incorporated herein by reference.

5 The term "lower alkyl" as used herein, and unless otherwise specified, refers to a C₁ to C₆ saturated straight, branched, or if appropriate, cyclic as in cyclopropyl, for eg., alkyl group, including both substituted and unsubstituted forms. Unless otherwise specifically stated in this application, when alkyl is a suitable moiety, lower alkyl is preferred. Similarly, when alkyl or lower alkyl is a suitable moiety, unsubstituted alkyl
10 or lower alkyl is preferred.

 The terms "alkylamino" and "arylamino" refer to an amino group that has one or two alkyl or aryl substituents, respectively.

 The term "protected" as used herein and, unless otherwise defined, refers to a group that is added to an oxygen, nitrogen or phosphorus atom to prevent its further
15 reaction or for other purposes. Numerous oxygen and nitrogen protecting groups are known to those skilled in the art of organic synthesis.

 The term "aryl" as used herein and, unless otherwise specified, refers to phenyl, biphenyl or naphthyl, and preferably phenyl. The term includes both substituted and unsubstituted moieties. The aryl group can be substituted with one or more moieties
20 selected from the group consisting of alkyl, hydroxyl, amino, alkylamino, arylamino, alkoxy, aryloxy, nitro, cyano, thio, alkylthio, carboxamido, carboxylate, sulfonic acid, sulfate, phosphonic acid, phosphate, or phosphonate, either unprotected or protected as necessary, as known to those skilled in the art, for eg., as taught in Greene et al., *Protective Groups in Organic Synthesis*, John Wiley and Sons, Second Edition (1991),
25 incorporated herein by reference.


 The terms "alkaryl" and "akylaryl" refer to an alkyl group with an aryl substituent.

 The terms "aralkyl" and "arylalkyl" refer to an aryl group with an alkyl substituent.

 The term "halo" as used herein includes bromo, chloro, iodo and fluoro.

5 The term purine base includes, but is not limited to, adenine, 2-azapurine bases that are optionally substituted imidazo-triazines, imidazo-pyridazines, pyrrolo-pyridazines, pyrrolo-triazines, triazolo-triazines including triazolo[4,5-*d*]triazines, pyrazolo-triazines including pyrazolo[4,5-*d*] triazines, N⁶-alkylpurines, N⁶-acylpurines (wherein acyl is C(O)(alkyl, aryl, alkylaryl, or arylalkyl), N⁶-benzylpurine, N⁶-halopurine, N⁶-vinylpurine, N⁶-acetylenic purine, N⁶-acyl purine, N⁶-hydroxyalkyl purine, N⁶-thioalkyl purine, N²-alkylpurines, N²-alkyl-6-thiopurines, C⁵-hydroxyalkyl purine, N²-alkylpurines, N²-alkyl-6-thiopurines, triazolopyridinyl, imidazolopyridinyl, pyrrolopyrimidinyl, and pyrazolopyrimidinyl.

10 The Base maybe selected from the group consisting of:



15 The term "acyl" refers to a carboxylic acid ester in which the non-carbonyl moiety of the ester group is selected from straight, branched, or cyclic alkyl or lower alkyl; alkoxyalkyl including methoxymethyl; aralkyl including benzyl; aryloxyalkyl such as phenoxymethyl; aryl including phenyl optionally substituted with halogen, C₁-C₆ alkyl

or C₁-C₆ alkoxy; sulfonate esters such as alkyl or aralkyl sulphonyl including methanesulfonyl; the mono-, di- or triphosphate ester; trityl or monomethoxytrityl; substituted benzyl; trialkylsilyl as, for eg., dimethyl-*t*-butylsilyl or diphenylmethylsilyl. Aryl groups in the esters optimally comprise a phenyl group. The term "lower acyl" refers to an acyl group in which the non-carbonyl moiety is lower alkyl.

As used herein, the terms "substantially free of" and "substantially in the absence of" refer to a nucleoside composition that includes at least 85-90% by weight, preferably 95%-98% by weight, and even more preferably 99%-100% by weight, of the designated enantiomer of that nucleoside. In a preferred embodiment, the compounds listed in the methods and compounds of this invention are substantially free of enantiomers other than for the one designated.

Similarly, the term "isolated" refers to a nucleoside composition that includes at least 85%-90% by weight, preferably 95%-98% by weight, and even more preferably 99%-100% by weight, of the nucleoside, the remainder comprising other chemical species or enantiomers.

The term "independently" is used herein to indicate that a variable is applied in any one instance without regard to the presence or absence of a variable having that same or a different definition within the same compound. Thus, in a compound in which R" appears twice and is defined as "independently carbon or nitrogen", both R"s can be carbon, both R"s can be nitrogen, or one R" can be carbon and the other nitrogen.

The term "host", as used herein, refers to a unicellular or multicellular organism in which the virus can replicate, including cell lines and animals, and preferably a human. Alternatively, the host can be carrying a part of the flavivirus or pestivirus genome, whose replication or function can be altered by the compounds of the present invention. The term host specifically refers to infected cells, cells transfected with all or part of the flavivirus or pestivirus genome and animals, in particular, primates (including chimpanzees) and humans. In most animal applications of the present invention, the host is a human patient. Veterinary applications, in certain indications, however, are clearly anticipated by the present invention such as in chimpanzees.

The term "pharmaceutically acceptable salt or prodrug" is used throughout the specification to describe any pharmaceutically acceptable form (ester, phosphate ester,

salt of an ester or a related group) of a nucleoside compound, which, upon administration to a patient, provides the nucleoside compound. Pharmaceutically acceptable salts include those derived from pharmaceutically acceptable inorganic or organic bases and acids. Suitable salts include those derived from alkali metals such as potassium and sodium, alkaline earth metals such as calcium and magnesium, among numerous other acids well known in the pharmaceutical art. Pharmaceutically acceptable prodrugs refer to a compound that is metabolized, for example, hydrolyzed or oxidized, in the host to form the compound of the present invention. Typical examples of prodrugs include compounds that have biologically labile protecting groups on a functional moiety of the active compound. Prodrugs include compounds that can be oxidized, reduced, aminated, deaminated, hydroxylated, dehydroxylated, hydrolyzed, dehydrolyzed, alkylated, dealkylated, acylated, deacylated, phosphorylated, dephosphorylated to produce the active compound. The compounds of this invention possess antiviral activity against flavivirus, pestivirus or HCV, or are metabolized to a compound that exhibits such activity.

Nucleoside Prodrug Formulations

Any of the nucleosides described herein can be administered as a nucleotide prodrug to increase the activity, bioavailability, stability or otherwise alter the properties of the nucleoside. A number of nucleotide prodrug ligands are known. In general, alkylation, acylation or other lipophilic modification of the mono-, di- or triphosphate of the nucleoside reduces polarity and allows passage into cells. Examples of substituent groups that can replace one or more hydrogens on the phosphate moiety are alkyl, aryl, steroids, carbohydrates, including sugars, 1,2-diacylglycerol, alcohols, acyl (including lower acyl); alkyl (including lower alkyl); sulfonate ester including alkyl or arylalkyl sulfonyl including methanesulfonyl and benzyl, wherein the phenyl group is optionally substituted with one or more substituents as provided in the definition of an aryl given herein; optionally substituted arylsulfonyl; a lipid, including a phospholipid; an amino acid residue or derivative; a carbohydrate; a peptide; cholesterol; or other pharmaceutically acceptable leaving group which, when administered *in vivo*, provides a compound wherein R^1 is independently H or phosphate. Many more are described in R.

Jones and N. Bischoferger, *Antiviral Research*, 1995, 27:1-17. Any of these can be used in combination with the disclosed nucleosides to achieve a desired effect.

5 In cases where compounds are sufficiently basic or acidic to form stable nontoxic acid or base salts, administration of the compound as a pharmaceutically acceptable salt may be appropriate. Examples of pharmaceutically acceptable salts are organic acid addition salts formed with acids, which form a physiological acceptable anion, for example, tosylate, methanesulfonate, acetate, citrate, malonate, tartarate, succinate, benzoate, ascorbate, α -ketoglutarate, and α -glycerophosphate. Suitable inorganic salts may also be formed, including, sulfate, nitrate, bicarbonate, and carbonate salts.

10 Pharmaceutically acceptable salts may be obtained using standard procedures well known in the art, for example by reacting a sufficiently basic compound such as an amine with a suitable acid affording a physiologically acceptable anion. Alkali metal (for example, sodium, potassium or lithium) or alkaline earth metal (for example calcium) salts of carboxylic acids can also be made.

15 The active nucleoside can also be provided as a 5'-phosphoether lipid or a 5'-ether lipid, as disclosed in the following references, which are incorporated by reference herein: Kucera, L.S., N. Iyer, E. Leake, A. Raen, Modest E.K., D.L.W., and C. Piantadosi. 1990. "Novel membrane-interactive ether lipid analogs that inhibit infectious HIV-1 production and induce defective virus formation." *AIDS Res. Hum. Retro Viruses*. 6:491-501; Piantadosi, C., J. Marasco C.J., S.L. Morris-Natschke, K.L. Meyer, F. Gumus, J.R. Surles, K.S. Ishaq, L.S. Kucera, N. Iyer, C.A. Wallen, S. Piantadosi, and E.J. Modest. 1991.

25 Nonlimiting examples of U.S. patents that disclose suitable lipophilic substituents that can be covalently incorporated into the nucleoside, preferably at the 5'-OH position of the nucleoside or lipophilic preparations, include U.S. Patent Nos. 5,149,794 (Sep. 22, 1992, Yatvin *et al.*); 5,194,654 (Mar. 16, 1993, Hostetler *et al.*, and 5,223,263 (June 29, 1993, Hostetler *et al.*); all of which are incorporated herein by reference. Foreign patent applications that disclose lipophilic substituents that can be attached to the nucleosides of the present invention, or lipophilic preparations, include WO 89/02733, WO 90/00555, 30 WO 91/16920, WO 91/18914, WO 93/00910, WO 94/26273, WO 96/15132, EP 0 350 287, EP 93917054.4, and WO 91/19721.

Combination and Alternation Therapy

It has been recognized that drug-resistant variants of HCV can emerge after prolonged treatment with an antiviral agent. Drug resistance most typically occurs by mutation of a gene that encodes for an enzyme used in viral replication. The efficacy of a drug against HCV infection can be prolonged, augmented, or restored by administering the compound in combination or alternation with a second, and perhaps third, antiviral compound that induces a different mutation from that caused by the principle drug. Alternatively, the pharmacokinetics, biodistribution or other parameter of the drug can be altered by such combination or alternation therapy. In general, combination therapy is typically preferred over alternation therapy because it induces multiple simultaneous stresses on the virus.

Any of the HCV treatments described in the Background of the Invention can be used in combination or alternation with the compounds described in this specification. Nonlimiting examples include:

(1) Interferon

Interferons (IFNs) are compounds that have been commercially available for the treatment of chronic hepatitis for nearly a decade. IFNs are glycoproteins produced by immune cells in response to viral infection. IFNs inhibit viral replication of many viruses, including HCV, and when used as the sole treatment for hepatitis C infection, IFN suppresses serum HCV-RNA to undetectable levels. Additionally, IFN normalizes serum amino transferase levels. Unfortunately, the effects of IFN are temporary and a sustained response occurs in only 8%-9% of patients chronically infected with HCV (Gary L. Davis. Gastroenterology 118:S104-S114, 2000).

A number of patents disclose HCV treatments using interferon-based therapies. For example, U.S. Patent No. 5,980,884 to Blatt et al. discloses methods for re-treatment of patients afflicted with HCV using consensus interferon. U.S. Patent No. 5,942,223 to Bazer et al. discloses an anti-HCV therapy using ovine or bovine interferon-tau. U.S. Patent No. 5,928,636 to Alber et al. discloses the combination therapy of interleukin-12 and interferon alpha for the treatment of infectious diseases including HCV. U.S. Patent No. 5,908,621 to Glue et al. discloses the use of polyethylene glycol modified interferon for the treatment of HCV. U.S. Patent No. 5,849,696 to Chretien et al. discloses the use

of thymosins, alone or in combination with interferon, for treating HCV. U.S. Patent No. 5,830,455 to Valtuena et al. discloses a combination HCV therapy employing interferon and a free radical scavenger. U.S. Patent No. 5,738,845 to Imakawa discloses the use of human interferon tau proteins for treating HCV. Other interferon-based treatments for HCV are disclosed in U.S. Patent No. 5,676,942 to Testa et al., U.S. Patent No. 5,372,808 to Blatt et al., and U.S. Patent No. 5,849,696.

(2) Ribavirin (Battaglia, A.M. et al., Ann. Pharmacother, 2000,. 34, 487-494); Berenguer, M. et al. Antivir. Ther., 1998, 3 (Suppl. 3), 125-136).

Ribavirin (1- β -D-ribofuranosyl-1-1,2,4-triazole-3-carboxamide) is a synthetic, non-interferon-inducing, broad spectrum antiviral nucleoside analog. It is sold under the trade names VirazoleTM (The Merck Index, 11th edition, Editor: Budavari, S., Merck & Co., Inc., Rahway, NJ, p1304, 1989); Rebetol (Schering Plough) and Co-Pegabus (Roche). United States Patent No. 3,798,209 and RE29,835 (ICN Pharmaceuticals) disclose and claim ribavirin. Ribavirin is structurally similar to guanosine, and has in vitro activity against several DNA and RNA viruses including Flaviviridae (Gary L. Davis. Gastroenterology 118:S104-S114, 2000). U.S. Patent No 4,211,771 (to ICN Pharmaceuticals) discloses the use of ribavirin as an antiviral agent. Ribavirin reduces serum amino transferase levels to normal in 40% of patients, but it does not lower serum levels of HCV-RNA (Gary L. Davis. Gastroenterology 118:S104-S114, 2000). Thus, ribavirin alone is not effective in reducing viral RNA levels. Additionally, ribavirin has significant toxicity and is known to induce anemia.

Combination of Interferon and Ribavirin

Schering-Plough sells ribavirin as Rebetol® capsules (200 mg) for administration to patients with HCV. The U.S. FDA has approved Rebetol capsules to treat chronic HCV infection in combination with Schering's alpha interferon-2b products Intron® A and PEG-IntronTM. Rebetol capsules are not approved for monotherapy (i.e., administration independent of Intron®A or PEG-Intron), although Intron A and PEG-Intron are approved for monotherapy (i.e., administration without ribavirin). Hoffman La Roche is selling ribavirin under the name Co-Pegabus in Europe and the United

States, also for use in combination with interferon for the treatment of HCV. Other alpha interferon products include Roferon-A (Hoffmann-La Roche), Infergen® (Intermune, formerly Amgen's product), and Wellferon® (Wellcome Foundation) are currently FDA-approved for HCV monotherapy. Interferon products currently in development for HCV include: Roferon-A (interferon alfa-2a) by Roche, PEGASYS (pegylated interferon alfa-2a) by Roche, INFERGEN (interferon alfacon-1) by InterMune, OMNIFERON (natural interferon) by Viragen, ALBUFERON by Human Genome Sciences, REBIF (interferon beta-1a) by Ares-Serono, Omega Interferon by BioMedicine, Oral Interferon Alpha by Amarillo Biosciences, and Interferon gamma-1b by InterMune.

The combination of IFN and ribavirin for the treatment of HCV infection has been reported to be effective in the treatment of IFN naïve patients (for example, Battaglia, A.M. et al., *Ann. Pharmacother.* 34:487-494, 2000). Combination treatment is effective both before hepatitis develops and when histological disease is present (for example, Berenguer, M. et al. *Antivir. Ther.* 3(Suppl. 3):125-136, 1998). Currently, the most effective therapy for HCV is combination therapy of pegylated interferon with ribavirin (2002 NIH Consensus Development Conference on the Management of Hepatitis C). However, the side effects of combination therapy can be significant and include hemolysis, flu-like symptoms, anemia, and fatigue (Gary L. Davis. *Gastroenterology* 118:S104-S114, 2000).

(3) Protease inhibitors have been developed for the treatment of *Flaviviridae* infections. Examples, include, but are not limited to the following

Substrate-based NS3 protease inhibitors (see, for example, Attwood *et al.*, *Antiviral peptide derivatives*, PCT WO 98/22496, 1998; Attwood *et al.*, *Antiviral Chemistry and Chemotherapy* 1999, 10, 259-273; Attwood *et al.*, *Preparation and use of amino acid derivatives as anti-viral agents*, German Patent Pub. DE 19914474; Tung *et al.* *Inhibitors of serine proteases, particularly hepatitis C virus NS3 protease*, PCT WO 98/17679), including alphaketoamides and hydrazinoureas, and inhibitors that terminate in an electrophile such as a boronic acid or phosphonate (see, for example, Llinas-Brunet *et al.*, *Hepatitis C inhibitor peptide analogues*, PCT WO 99/07734);

Non-substrate-based inhibitors such as 2,4,6-trihydroxy-3-nitro-benzamide derivatives (see, for example, Sudo K. *et al.*, *Biochemical and Biophysical Research*

Communications, 1997, 238, 643-647; Sudo K. *et al.* *Antiviral Chemistry and Chemotherapy*, 1998, 9, 186), including RD3-4082 and RD3-4078, the former substituted on the amide with a 14 carbon chain and the latter processing a *para*-phenoxyphenyl group;

5 Phenanthrenequinones possessing activity against protease, for example in a SDS-PAGE and/or autoradiography assay, such as, for example, Sch 68631, isolated from the fermentation culture broth of *Streptomyces* sp., (see, for example, Chu M. *et al.*, *Tetrahedron Letters*, 1996, 37, 7229-7232), and Sch 351633, isolated from the fungus *Penicillium griseofulvum*, which demonstrates activity in a scintillation proximity assay
10 (see, for example, Chu M. *et al.*, *Bioorganic and Medicinal Chemistry Letters* 9, 1949-1952); and

Selective NS3 inhibitors, for example, based on the macromolecule elgin c, isolated from leech (see, for example, Qasim M.A. *et al.*, *Biochemistry*, 1997, 36, 1598-1607). Nanomolar potency against the HCV NS3 protease enzyme has been achieved by
15 the design of selective inhibitors based on the macromolecule eglin c. Eglin c, isolated from leech, is a potent inhibitor of several serine proteases such as *S. griseus* proteases A and B, α -chymotrypsin, chymase and subtilisin.

Several U.S. patents disclose protease inhibitors for the treatment of HCV. Non-limiting examples include, but are not limited to the following. U.S. Patent No.
20 6,004,933 to Spruce *et al.* discloses a class of cysteine protease inhibitors for inhibiting HCV endopeptidase. U.S. Patent No. 5,990,276 to Zhang *et al.* discloses synthetic inhibitors of hepatitis C virus NS3 protease. The inhibitor is a subsequence of a substrate of the NS3 protease or a substrate of the NS4A cofactor. The use of restriction enzymes to treat HCV is disclosed in U.S. Patent No. 5,538,865 to Reyes *et al.* Peptides
25 as NS3 serine protease inhibitors of HCV are disclosed in WO 02/008251 to Corvas International, Inc, and WO 02/08187 and WO 02/008256 to Schering Corporation. HCV inhibitor tripeptides are disclosed in US Patent Nos. 6,534,523, 6,410,531, and 6,420,380 to Boehringer Ingelheim and WO 02/060926 to Bristol Myers Squibb. Diaryl peptides as NS3 serine protease inhibitors of HCV are disclosed in WO 02/48172 to Schering
30 Corporation. Imidazoleidinones as NS3 serine protease inhibitors of HCV are disclosed in WO 02/08198 to Schering Corporation and WO 02/48157 to Bristol Myers Squibb.

WO 98/17679 to Vertex Pharmaceuticals and WO 02/48116 to Bristol Myers Squibb also disclose HCV protease inhibitors.

5 (4) Thiazolidine derivatives, for example, that show relevant inhibition in a reverse-phase HPLC assay with an NS3/4A fusion protein and NS5A/5B substrate (see, for example, Sudo K. *et al.*, *Antiviral Research*, **1996**, 32, 9-18), especially compound RD-1-6250, possessing a fused cinnamoyl moiety substituted with a long alkyl chain, RD4 6205 and RD4 6193;

10 (5) Thiazolidines and benzanilides, for example, as identified in Kakiuchi N. *et al.* *J. EBS Letters* 421, 217-220; Takeshita N. *et al.* *Analytical Biochemistry*, **1997**, 247, 242-246;

15 (6) Helicase inhibitors (see, for example, Diana G.D. *et al.*, *Compounds, compositions and methods for treatment of hepatitis C*, U.S. Pat. No. 5,633,358; Diana G.D. *et al.*, *Piperidine derivatives, pharmaceutical compositions thereof and their use in the treatment of hepatitis C*, PCT WO 97/36554);

(7) Polymerase inhibitors such as

i) nucleotide analogues, such as gliotoxin (see, for example, Ferrari R. *et al.* *Journal of Virology*, **1999**, 73, 1649-1654);

20 ii) the natural product cerulenin (see, for example, Lohmann V. *et al.*, *Virology*, **1998**, 249, 108-118); and

25 iii) non-nucleoside polymerase inhibitors, including, for example, compound R803 (see, for example, WO 04/018463 A2 and WO 03/040112 A1, both to Rigel Pharmaceuticals, Inc.); substituted diamine pyrimidines (see, for example, WO 03/063794 A2 to Rigel Pharmaceuticals, Inc.); benzimidazole derivatives (see, for example, *Bioorg. Med. Chem. Lett.*, **2004**, 14:119-124 and *Bioorg. Med. Chem. Lett.*, **2004**, 14:967-971, both to Boehringer Ingelheim Corporation); N,N-disubstituted phenylalanines (see, for example, *J. Biol. Chem.*, **2003**, 278:9495-98 and *J. Med. Chem.*, **2003**, 13:1283-85, both to

30

5 Shire Biochem, Inc.); substituted thiophene-2-carboxylic acids (see, for example, *Bioorg. Med. Chem. Lett.*, **2004**, *14*:793-796 and *Bioorg. Med. Chem. Lett.*, **2004**, *14*:797-800, both to Shire Biochem, Inc.); α,γ -diketoacids (see, for example, *J. Med. Chem.*, **2004**, *14-17* and WO 00/006529 A1, both to Merck & Co., Inc.); and meconic acid derivatives (see, for example, *Bioorg. Med. Chem. Lett.*, **2004**, 3257-3261, WO 02/006246 A1 and WO03/062211 A1, all to IRBM Merck & Co., Inc.);

10 (8) Antisense phosphorothioate oligodeoxynucleotides (S-ODN) complementary, for example, to sequence stretches in the 5' non-coding region (NCR) of the virus (see, for example, Alt M. *et al.*, *Hepatology*, **1995**, *22*, 707-717), or to nucleotides 326-348 comprising the 3' end of the NCR and nucleotides 371-388 located in the core coding region of the HCV RNA (see, for example, Alt M. *et al.*, *Archives of Virology*, **1997**, *142*, 589-599; Galderisi U. *et al.*, *Journal of Cellular Physiology*, **1999**, *181*, 251-257).

15 (9) Inhibitors of IRES-dependent translation (see, for example, Ikeda N *et al.*, *Agent for the prevention and treatment of hepatitis C*, Japanese Patent Pub. JP-08268890; Kai Y. *et al.* *Prevention and treatment of viral diseases*, Japanese Patent Pub. JP-10101591).

20 (10) Nuclease-resistant ribozymes (see, for example, Maccjak, D. J. *et al.*, *Hepatology* **1999**, *30*, abstract 995; U.S. Patent No. 6,043,077 to Barber *et al.*, and U.S. Patent Nos. 5,869,253 and 5,610,054 to Draper *et al.*).

(11) Nucleoside analogs have also been developed for the treatment of Flaviviridae infections.

25 Idenix Pharmaceuticals, Ltd. discloses branched nucleosides, and their use in the treatment of HCV and flaviviruses and pestiviruses in US Patent Publication Nos. 2003/0050229 A1, 2004/0097461 A1, 2004/0101535 A1, 2003/0060400 A1, 2004/0102414 A1, 2004/0097462 A1, and 2004/0063622 A1 which correspond to International Publication Nos. WO 01/90121 and WO 01/92282. A method for the treatment of hepatitis C infection (and flaviviruses and pestiviruses) in humans and other host animals is disclosed in the Idenix publications that includes administering an effective amount of a biologically active 1', 2', 3' or 4'-branched β -D or β -L nucleosides
30 or a pharmaceutically acceptable salt or prodrug thereof, administered either alone or in

combination, optionally in a pharmaceutically acceptable carrier. See also U.S. Patent Publication Nos. 2004/0006002 and 2004/0006007 as well as WO 03/026589 and WO 03/026675. Idenix Pharmaceuticals, Ltd. also discloses in US Patent Publication No. 2004/0077587 pharmaceutically acceptable branched nucleoside prodrugs, and their use
5 in the treatment of HCV and flaviviruses and pestiviruses in prodrugs. See also PCT Publication Nos. WO 04/002422, WO 04/002999, and WO 04/003000. Further, Idenix Pharmaceuticals, Ltd. also discloses in WO 04/046331 Flaviviridae mutations caused by biologically active 2'-branched β -D or β -L nucleosides or a pharmaceutically acceptable salt or prodrug thereof.

10 Biota Inc. discloses various phosphate derivatives of nucleosides, including 1', 2', 3' or 4'-branched β -D or β -L nucleosides, for the treatment of hepatitis C infection in International Patent Publication WO 03/072757.

Emory University and the University of Georgia Research Foundation, Inc. (UGARF) discloses the use of 2'-fluoronucleosides for the treatment of HCV in US
15 Patent No. 6,348,587. See also US Patent Publication No. 2002/0198171 and International Patent Publication WO 99/43691.

BioChem Pharma Inc. (now Shire Biochem, Inc.) discloses the use of various 1,3-dioxolane nucleosides for the treatment of a *Flaviviridae* infection in US Patent No. 6,566,365. See also US Patent Nos. 6,340,690 and 6,605,614; US Patent Publication
20 Nos. 2002/0099072 and 2003/0225037, as well as International Publication No. WO 01/32153 and WO 00/50424..

BioChem Pharma Inc. (now Shire Biochem, Inc.) also discloses various other 2'-halo, 2'-hydroxy and 2'-alkoxy nucleosides for the treatment of a *Flaviviridae* infection in US Patent Publication No. 2002/0019363 as well as International Publication No. WO
25 01/60315 (PCT/CA01/00197; filed February 19, 2001).

ICN Pharmaceuticals, Inc. discloses various nucleoside analogs that are useful in modulating immune response in US Patent Nos. 6,495,677 and 6,573,248. See also WO 98/16184, WO 01/68663, and WO 02/03997.

US Patent No. 6,660,721; US Patent Publication Nos. 2003/083307 A1,
30 2003/008841 A1, and 2004/0110718; as well as International Patent Publication Nos. WO 02/18404; WO 02/100415, WO 02/094289, and WO 04/043159; filed by F.

Hoffmann-La Roche AG, discloses various nucleoside analogs for the treatment of HCV RNA replication.

Pharmasset Limited discloses various nucleosides and antimetabolites for the treatment of a variety of viruses, including *Flaviviridae*, and in particular HCV, in US
5 Patent Publication Nos. 2003/0087873, 2004/0067877, 2004/0082574, 2004/0067877, 2004/002479, 2003/0225029, and 2002/00555483, as well as International Patent Publication Nos. WO 02/32920, WO 01/79246, WO 02/48165, WO 03/068162, WO 03/068164 and WO 2004/013298.

Merck & Co., Inc. and Isis Pharmaceuticals disclose in US Patent Publication No.
10 2002/0147160, 2004/0072788, 2004/0067901, and 2004/0110717; as well as the corresponding International Patent Publication Nos. WO 02/057425 (PCT/US02/01531; filed January 18, 2002) and WO 02/057287 (PCT/US02/03086; filed January 18, 2002) various nucleosides, and in particular several pyrrolopyrimidine nucleosides, for the treatment of viruses whose replication is dependent upon RNA-dependent RNA
15 polymerase, including *Flaviviridae*, and in particular HCV. See also WO 2004/000858, WO 2004/003138, WO 2004/007512, and WO 2004/009020.

US Patent Publication No. 2003/028013 A1 as well as International Patent Publication Nos. WO 03/051899, WO 03/061576, WO 03/062255 WO 03/062256, WO 03/062257, and WO 03/061385, filed by Ribapharm, also are directed to the use of
20 certain nucleoside analogs to treat hepatitis C virus.

Genelabs Technologies disclose in US Patent Publication No. 2004/0063658 as well as International Patent Publication Nos. WO 03/093290 and WO 04/028481 various base modified derivatives of nucleosides, including 1', 2', 3' or 4'-branched β -D or β -L nucleosides, for the treatment of hepatitis C infection.

25 Eldrup et al. (Oral Session V, Hepatitis C Virus, *Flaviviridae*; 16th International Conference on Antiviral Research (April 27, 2003, Savannah, Ga.) p. A75) described the structure activity relationship of 2'-modified nucleosides for inhibition of HCV.

Bhat et al (Oral Session V, Hepatitis C Virus, *Flaviviridae*; 16th International Conference on Antiviral Research (April 27, 2003, Savannah, Ga.); p A75) describe the
30 synthesis and pharmacokinetic properties of nucleoside analogues as possible inhibitors

of HCV RNA replication. The authors report that 2'-modified nucleosides demonstrate potent inhibitory activity in cell-based replicon assays.

Olsen et al. (Oral Session V, Hepatitis C Virus, *Flaviviridae*; 16th International Conference on Antiviral Research (April 27, 2003, Savannah, Ga.) p A76) also described the effects of the 2'-modified nucleosides on HCV RNA replication.

(12) Other miscellaneous compounds including 1-amino-alkylcyclohexanes (for example, U.S. Patent No. 6,034,134 to Gold *et al.*), alkyl lipids (for example, U.S. Pat. No. 5,922,757 to Chojkier *et al.*), vitamin E and other antioxidants (for example, U.S. Pat. No. 5,922,757 to Chojkier *et al.*), squalene, amantadine, bile acids (for example, U.S. Pat. No. 5,846,964 to Ozeki *et al.*), N-(phosphonoacetyl)-L-aspartic acid (for example, U.S. Pat. No. 5,830,905 to Diana *et al.*), benzenedicarboxamides (for example, U.S. Pat. No. 5,633,388 to Diana *et al.*), polyadenylic acid derivatives (for example, U.S. Pat. No. 5,496,546 to Wang *et al.*), 2',3'-dideoxyinosine (for example, U.S. Pat. No. 5,026,687 to Yarchoan *et al.*), benzimidazoles (for example, U.S. Pat. No. 5,891,874 to Colacino *et al.*), plant extracts (for example, U.S. Patent No. 5,837,257 to Tsai *et al.*, U.S. Patent No. 5,725,859 to Omer *et al.*, and U.S. Patent No. 6,056,961), and piperidenes (for example, U.S. Patent No. 5,830,905 to Diana *et al.*).

(13) Other compounds currently in clinical development for treatment of hepatitis C virus include, for example: Interleukin-10 by Schering-Plough, IP-501 by Interneuron, Merimebodib VX-497 by Vertex, AMANTADINE® (Symmetrel) by Endo Labs Solvay, HEPTAZYME® by RPI, IDN-6556 by Idun Pharma., XTL-002 by XTL., HCV/MF59 by Chiron, CIVACIR® (Hepatitis C Immune Globulin) by NABI, LEVOVIRIN® by ICN/Ribapharm, VIRAMIDINE® by ICN/Ribapharm, ZADAXIN® (thymosin alfa-1) by Sci Clone, thymosin plus pegylated interferon by Sci Clone, CEPLENE® (histamine dihydrochloride) by Maxim, VX 950 / LY 570310 by Vertex/Eli Lilly, ISIS 14803 by Isis Pharmaceutical/Elan, IDN-6556 by Idun Pharmaceuticals, Inc., JTK 003 by AKROS Pharma, BILN-2061 by Boehringer Ingelheim, CellCept (mycophenolate mofetil) by Roche, T67, a β -tubulin inhibitor, by Tularik, a therapeutic vaccine directed to E2 by Innogenetics, FK788 by Fujisawa Healthcare, Inc., IdB 1016 (Siliphos, oral silybin-phosphatidylcholine phytosome), RNA replication inhibitors (VP50406) by

ViroPharma/Wyeth, therapeutic vaccine by Intercell, therapeutic vaccine by Epimmune/Genencor, IRES inhibitor by Anadys, ANA 245 and ANA 246 by Anadys, immunotherapy (Therapore) by Avant, protease inhibitor by Corvas/SChering, helicase inhibitor by Vertex, fusion inhibitor by Trimeris, T cell therapy by CellExSys, polymerase inhibitor by Biocryst, targeted RNA chemistry by PTC Therapeutics, Dication by Immtech, Int., protease inhibitor by Agouron, protease inhibitor by Chiron/Medivir, antisense therapy by AVI BioPharma, antisense therapy by Hybridon, hemopurifier by Aethlon Medical, therapeutic vaccine by Merix, protease inhibitor by Bristol-Myers Squibb/Axys, Chron-VacC, a therapeutic vaccine, by Tripep, UT 231B by United Therapeutics, protease, helicase and polymerase inhibitors by Genelabs Technologies, IRES inhibitors by Immusol, R803 by Rigel Pharmaceuticals, INFERGEN® (interferon alphacon-1) by InterMune, OMNIFERON® (natural interferon) by Viragen, ALBUFERON® by Human Genome Sciences, REBIF® (interferon beta-1a) by Ares-Serono, Omega Interferon by BioMedicine, Oral Interferon Alpha by Amarillo Biosciences, interferon gamma, interferon tau, and Interferon gamma- 1b by InterMune.

Pharmaceutical Compositions

Hosts, including humans, infected with pestivirus, flavivirus, HCV or another organism replicating through a RNA-dependent RNA viral polymerase, can be treated by administering to the patient an effective amount of the active compound or a pharmaceutically acceptable prodrug or salt thereof in the presence of a pharmaceutically acceptable carrier or diluent. The active materials can be administered by any appropriate route, for example, orally, parenterally, intravenously, intradermally, subcutaneously, or topically, in liquid or solid form.

A preferred dose of the compound for pestivirus, flavivirus or HCV will be in the range from about 1 to 50 mg/kg, preferably 1 to 20 mg/kg, of body weight per day, more generally 0.1 to about 100 mg per kilogram body weight of the recipient per day. The effective dosage range of the pharmaceutically acceptable salts and prodrugs can be calculated based on the weight of the parent nucleoside to be delivered. If the salt or

prodrug exhibits activity in itself, the effective dosage can be estimated as above using the weight of the salt or prodrug, or by other means known to those skilled in the art.

5 The compound is conveniently administered in unit any suitable dosage form, including but not limited to one containing 7 to 3000 mg, or 70 to 1400 mg of active ingredient per unit dosage form. An oral dosage in one embodiment is 50-1000 mg. In another embodiment, the dosage form contains 0.5-500 mg; or 0.5-100 mg; or 0.5-50 mg; or 0.5-25 mg; or 1.0-10 mg.

10 Ideally the active ingredient should be administered to achieve peak plasma concentrations of the active compound of from about 0.2 to 70 μ M, preferably about 1.0 to 10 μ M. This may be achieved, for example, by the intravenous injection of a 0.1 to 5% solution of the active ingredient, optionally in saline, or administered as a bolus of the active ingredient.

15 The concentration of active compound in the drug composition will depend on absorption, inactivation and excretion rates of the drug as well as other factors known to those of skill in the art. It is to be noted that dosage values will also vary with the severity of the condition to be alleviated. It is to be further understood that for any particular subject, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions, and that the concentration ranges set
20 forth herein are exemplary only and are not intended to limit the scope or practice of the claimed composition. The active ingredient may be administered at once, or may be divided into a number of smaller doses to be administered at varying intervals of time.

25 A preferred mode of administration of the active compound is oral. Oral compositions will generally include an inert diluent or an edible carrier. They may be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition.

30 The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a

disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring. When the dosage unit form is a capsule, it can contain, in addition to material of the above type, a liquid carrier such as a fatty oil. In addition, dosage unit forms can contain various other materials which modify the physical form of the dosage unit, for example, coatings of sugar, shellac, or other enteric agents.

The compound can be administered as a component of an elixir, suspension, syrup, wafer, chewing gum or the like. A syrup may contain, in addition to the active compounds, sucrose as a sweetening agent and certain preservatives, dyes and colorings and flavors.

The compound or a pharmaceutically acceptable prodrug or salts thereof can also be mixed with other active materials that do not impair the desired action, or with materials that supplement the desired action, such as antibiotics, antifungals, anti-inflammatories, or other antivirals, including other nucleoside compounds. Solutions or suspensions used for parenteral, intradermal, subcutaneous, or topical application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. The parental preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

If administered intravenously, preferred carriers are physiological saline or phosphate buffered saline (PBS).

In a preferred embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters and polylactic acid.

Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation.


Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) are also preferred as pharmaceutically acceptable carriers. These may be prepared according to methods known to those skilled in the art, for example, as described in U.S. Patent No. 4,522,811 (which is incorporated herein by reference in its entirety). For example, liposome formulations may be prepared by dissolving appropriate lipid(s) (such as stearyl phosphatidyl ethanolamine, stearyl phosphatidyl choline, arachidoyl phosphatidyl choline, and cholesterol) in an inorganic solvent that is then evaporated, leaving behind a thin film of dried lipid on the surface of the container. An aqueous solution of the active compound or its monophosphate, diphosphate, and/or triphosphate derivatives is then introduced into the container. The container is then swirled by hand to free lipid material from the sides of the container and to disperse lipid aggregates, thereby forming the liposomal suspension.

Processes for the Preparation of Active Compounds

The nucleosides of the present invention can be synthesized by any means known in the art. In particular, the synthesis of the present nucleosides can be achieved by either alkylating the appropriately modified sugar, followed by glycosylation or glycosylation followed by alkylation of the nucleoside, though preferably alkylating the appropriately modified sugar, followed by glycosylation. The following non-limiting embodiments illustrate some general methodology to obtain the nucleosides of the present invention.

A. General Synthesis of 1'-C-branched Nucleosides

1'-C branched ribonucleosides of the following structures:



wherein

R is H, phosphate (including mono-, di-, or triphosphate or a stabilized phosphate
5 prodrug) or phosphonate;

n is 0-2;

when X is CH₂, CHOH, CH-alkyl, CH-alkenyl, CH-alkynyl, C-dialkyl, CH-O-alkyl, CH-
O-alkenyl, CH-O-alkynyl, CH-S-alkyl, CH-S-alkenyl, CH-S-alkynyl, CH-halogen, or
C-(halogen)₂,

10 then each R¹ and R^{1'} is independently H, OH, optionally substituted alkyl including
lower alkyl, azido, cyano, optionally substituted alkenyl or alkynyl, -C(O)O-(alkyl),
-C(O)O(lower alkyl), -C(O)O-(alkenyl), -C(O)O-(alkynyl), -O(acyl), -O(lower acyl),
-O(alkyl), -O(lower alkyl), -O(alkenyl), -O(alkynyl), halogen, halogenated alkyl,
-NO₂, -NH₂, -NH(lower alkyl), -N(lower alkyl)₂, -NH(acyl), -N(acyl)₂, -C(O)NH₂,
15 -C(O)NH(alkyl), -C(O)N(alkyl)₂, S(O)N-alkyl, S(O)N-alkenyl, S(O)N-alkynyl,
SCH-halogen, wherein alkyl, alkenyl, and/or alkynyl maybe optionally substituted;


when X is O, S[O]_n, NH, N-alkyl, N-alkenyl, N-alkynyl, S(O)N-alkyl, S(O)N-alkenyl,
S(O)N-alkynyl, or SCH-halogen,

20 then each R¹ and R^{1'} is independently H, optionally substituted alkyl including lower
alkyl, azido, cyano, optionally substituted alkenyl or alkynyl, -C(O)O-(alkyl),
-C(O)O(lower alkyl), -C(O)O-(alkenyl), -C(O)O-(alkynyl), halogenated alkyl,
-C(O)NH₂, -C(O)NH(alkyl), -C(O)N(alkyl)₂, -C(H)=N-NH₂, C(S)NH₂,
C(S)NH(alkyl), or C(S)N(alkyl)₂, wherein alkyl, alkenyl, and/or alkynyl maybe
optionally substituted;

25 when X* is CY³;

- then each R¹ is independently H, OH, optionally substituted alkyl including lower alkyl, azido, cyano, optionally substituted alkenyl or alkynyl, -C(O)O-(alkyl), -C(O)O(lower alkyl), -C(O)O-(alkenyl), -C(O)O-(alkynyl), -O(acyl), -O(lower acyl), -O(alkyl), -O(lower alkyl), -O(alkenyl), -O(alkynyl), halogen, halogenated alkyl, -NO₂, -NH₂, -NH(lower alkyl), -N(lower alkyl)₂, -NH(acyl), -N(acyl)₂, -C(O)NH₂, -C(O)NH(alkyl), and -C(O)N(alkyl)₂, wherein an optional substitution on alkyl, alkenyl, and/or alkynyl may be one or more halogen, hydroxy, alkoxy or alkylthio groups taken in any combination; and
- Y³ is hydrogen, alkyl, bromo, chloro, fluoro, iodo, azido, cyano, alkenyl, alkynyl, -C(O)O(alkyl), -C(O)O(lower alkyl), CF₃, -CONH₂, -CONH(alkyl), -CON(alkyl)₂;
- each R² and R³ independently is OH, NH₂, SH, F, Cl, Br, I, CN, NO₂, -C(O)NH₂, -C(O)NH(alkyl), -C(O)N(alkyl)₂, N₃, optionally substituted alkyl including lower alkyl, optionally substituted alkenyl or alkynyl, halogenated alkyl, -C(O)O-(alkyl), -C(O)O(lower alkyl), -C(O)O-(alkenyl), -C(O)O-(alkynyl), -O(acyl), -O(alkyl), -O(alkenyl), -O(alkynyl), -OC(O)NH₂, NC, C(O)OH, SCN, OCN, -S(alkyl), -S(alkenyl), -S(alkynyl), -NH(alkyl), -N(alkyl)₂, -NH(alkenyl), -NH(alkynyl), an amino acid residue or derivative, a prodrug or leaving group that provides OH *in vivo*, or an optionally substituted 3-7 membered heterocyclic ring having O, S and/or N independently as a heteroatom taken alone or in combination;
- each R^{2'} and R^{3'} independently is H; optionally substituted alkyl, alkenyl, or alkynyl; -C(O)O(alkyl), -C(O)O(lower alkyl), -C(O)O(alkenyl), -C(O)O(alkynyl), -C(O)NH₂, -C(O)NH(alkyl), -C(O)N(alkyl)₂, -O(acyl), -O(lower acyl), -O(alkyl), -O(lower alkyl), -O(alkenyl), halogen, halogenated alkyl and particularly CF₃, azido, cyano, NO₂, -S(alkyl), -S(alkenyl), -S(alkynyl), NH₂, -NH(alkyl), -N(alkyl)₂, -NH(alkenyl), -NH(alkynyl), -NH(acyl), or -N(acyl)₂, and R₃ at 3'-C may also be OH;

Base is selected from the group consisting of:



wherein

5 each A independently is N or C-R⁵;

W is H, OH, -O(acyl), -O(C₁₋₄ alkyl), -O(alkenyl), -O(alkynyl), -OC(O)R⁴R⁴, -OC(O)N R⁴R⁴, SH, -S(acyl), -S(C₁₋₄ alkyl), NH₂, NH(acyl), N(acyl)₂, NH(C₁₋₄ alkyl), N(C₁₋₄ alkyl)₂, -N(cycloalkyl) C₁₋₄ alkylamino, di(C₁₋₄ alkyl)amino, C₃₋₆ cycloalkylamino, halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, CN, SCN, OCN, SH, N₃, NO₂, NH=NH₂, N₃,
 10 NHOH, -C(O)NH₂, -C(O)NH(acyl), -C(O)N(acyl)₂, -C(O)NH(C₁₋₄ alkyl), -C(O)N(C₁₋₄ alkyl)₂, -C(O)N(alkyl)(acyl), or halogenated alkyl;

Z is O, S, NH, N-OH, N-NH₂, NH(alkyl), N(alkyl)₂, N-cycloalkyl, alkoxy, CN, SCN, OCN, SH, NO₂, NH₂, N₃, NH=NH, NH(alkyl), N(alkyl)₂, CONH₂, CONH(alkyl), or CON(alkyl)₂.

15 each R⁴ is independently H, acyl, or C₁₋₆ alkyl;

each R⁵ is independently H, Cl, Br, F, I, CN, OH, optionally substituted alkyl, alkenyl or alkynyl, carboxy, C(=NH)NH₂, C₁₋₄ alkoxy, C₁₋₄ alkyloxycarbonyl, N₃, NH₂, NH(alkyl), N(alkyl)₂, NO₂, N₃, halogenated alkyl especially CF₃, C₁₋₄ alkylamino, di(C₁₋₄ alkyl)amino, C₃₋₆ cycloalkylamino, C₁₋₆ alkoxy, SH, -S(C₁₋₄ alkyl), -S(C₁₋₄ alkenyl), -S(C₁₋₄ alkynyl), C₁₋₆ alkylthio, C₁₋₆ alkylsulfonyl, (C₁₋₄ alkyl)₀₋₂ aminomethyl, C₃₋₆ cycloalkylamino -alkenyl, -alkynyl, -(O)alkyl, -(O)alkenyl, -(O)alkynyl, -(O)acyl, -O(C₁₋₄ alkyl), -O(C₁₋₄ alkenyl), -O(C₁₋₄ alkynyl), -O-C(O)NH₂, -OC(O)N(alkyl), -OC(O)R'R'', -C(O)OH, C(O)O-alkyl, C(O)O-alkenyl, C(O)O-alkynyl, S-alkyl, S-acyl, S-alkenyl, S-alkynyl, SCN, OCN, NC, -C(O)-NH₂, C(O)NH(alkyl), C(O)N(alkyl)₂, C(O)NH(acyl), C(O)N(acyl)₂, (S)-NH₂, NH-alkyl, N(dialkyl)₂, NH-acyl, N-diacyl, or a 3-7 membered heterocycle having O, S, or N taken independently in any combination;

each R' and R'' independently is H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, halogen, halogenated alkyl, OH, CN, N₃, carboxy, C₁₋₄alkoxycarbonyl, NH₂, C₁₋₄ alkylamino, di(C₁₋₄ alkyl)amino, C₁₋₆ alkoxy, C₁₋₆ alkylsulfonyl, or (C₁₋₄ alkyl)₀₋₂ aminomethyl; and all tautomeric, enantiomeric and stereoisomeric forms thereof;

with the caveat that when X is S in Formula (I), then the compound is not 5-(4-aminoimidazo [4,5-d][1,2,3]triazin-7-yl)-2-hydroxymethyl-tetrahydro-thiophen-3-ol or 7-(4-hydroxy-5-hydroxy-methyl-tetrahydro-thiophen-2-yl)-3,7-dihydro-imidazo[4,5-d][1,2,3]triazin-4-one, can be prepared according to Schemes 1, 2 or 7 below.

Modification from the Lactone

The key starting material for this process is an appropriately substituted lactone. The lactone may be purchased or can be prepared by any known means including standard epimerization, substitution and cyclization techniques. The lactone optionally can be protected with a suitable protecting group, preferably with an acyl or silyl group, by methods well known to those skilled in the art, as taught by Greene et al., Protective Groups in Organic Synthesis, John Wiley and Sons, Second Edition, 1991. The protected lactone can then be coupled with a suitable coupling agent, such as an organometallic carbon nucleophile like a Grignard reagent, an organolithium, lithium


dialkylcopper or R^6-SiMe_3 in TAF with the appropriate non-protic solvent at a suitable temperature, to give the 1'-alkylated sugar.

The optionally activated sugar can then be coupled to the base by methods well known to those skilled in the art, as taught by Townsend, Chemistry of Nucleotides, Plenum Press, 1994. For example, an acylated sugar can be coupled to a silylated base with a Lewis acid such as tin tetrachloride, titanium tetrachloride, or trimethylsilyltriflate in the appropriate solvent at a suitable temperature.

Subsequently, the nucleoside can be deprotected by methods well known to those skilled in the art, as taught by Greene et al., Protective Groups in Organic Synthesis, John Wiley and Sons, Second Edition, 1991.

In a particular embodiment, the 1'-C-branched ribonucleoside is desired. The synthesis of a ribonucleoside is shown in Scheme 1. Alternatively, deoxyribonucleoside is desired. To obtain these nucleosides, the formed ribonucleoside can optionally be protected by methods well known to those skilled in the art, as taught by Greene et al., Protective Groups in Organic Synthesis, John Wiley and Sons, Second Edition, 1991, and then the 2'-OH can be reduced with a suitable reducing agent. Optionally, the 2'-OH can be activated to facilitate reduction as, for example, via the Barton reduction.

Scheme 1



Alternative Method for the Preparation of 1'-C-branched Nucleosides

The key starting material for this process is an appropriately substituted hexose. The hexose can be purchased or can be prepared by any known means including standard epimerization (as, for example, via alkaline treatment), substitution and coupling techniques. The hexose can be protected selectively to give the appropriate hexa-furanose, as taught by Townsend, Chemistry of Nucleosides and Nucleotides, Plenum Press, 1994.


The 1'-OH optionally can be activated to a suitable leaving group such as an acyl group or a halogen via acylation or halogenation, respectively. The optionally activated sugar can then be coupled to the base by methods well known to those skilled in the art, as taught by Townsend, Chemistry of Nucleosides and Nucleotides, Plenum Press, 1994. For example, an acylated sugar can be coupled to a silylated base with a Lewis acid, such as tin tetrachloride, titanium tetrachloride, or trimethylsilyltriflate in an appropriate

solvent at a suitable temperature. Alternatively, a halo-sugar can be coupled to a silylated base in the presence of trimethylsilyltriflate.

5 The 1'-CH₂-OH, if protected, selectively can be deprotected by methods well known in the art. The resultant primary hydroxyl can be reduced to give the methyl, using a suitable reducing agent. Alternatively, the hydroxyl can be activated prior to reduction to facilitate the reaction, i.e., via the Barton reduction. In an alternate embodiment, the primary hydroxyl can be oxidized to the aldehyde, then coupled with a carbon nucleophile such as a Grignard reagent, an organolithium, lithium dialkylcopper or R⁶-SiMe₃ in TAF with an appropriate non-protic solvent at a suitable temperature.

10 In a particular embodiment, the 1'-C-branched ribonucleoside is desired. The synthesis of a ribonucleoside is shown in Scheme 2. Alternatively, deoxyribonucleoside is desired. To obtain these nucleosides, the formed ribonucleoside optionally can be protected by methods well known to those skilled in the art, as taught by Greene et al., Protective Groups in Organic Synthesis, John Wiley and Sons, Second Edition, 1991,
15 and then the 2'-OH can be reduced with a suitable reducing agent. Optionally, the 2'-OH can be activated to facilitate reduction as, for example, via the Barton reduction.


Scheme 2



In addition, the L-enantiomers corresponding to the compounds of the invention can be prepared following the same general methods (1 or 2), beginning with the corresponding L-sugar or nucleoside L-enantiomer as the starting material.

5 General Synthesis of 2'-C-branched Nucleosides

2'-C-branched ribonucleosides of the following structures:



wherein R, R¹, R^{1'}, R², R^{2'}, R³, R^{3'}, X, X*, and Base are all as described above, can be prepared according to Schemes 3 or 4 below.

Glycosylation of the nucleobase with an appropriately modified sugar

5 The key starting material for this process is an appropriately substituted sugar with a 2'-OH and 2'-H, with an appropriate leaving group (LG), such as an acyl or halogen group, for example. The sugar can be purchased or can be prepared by any known means including standard epimerization, substitution, oxidation and/or reduction techniques. The substituted sugar can then be oxidized with an appropriate oxidizing agent in a compatible solvent at a suitable temperature to yield the 2'-modified sugar. Possible oxidizing agents are Jones' reagent (a mixture of chromic and sulfuric acids), Collins' reagent (dipyridine Cr(VI)oxide), Corey's reagent (pyridinium chlorochromate), pyridinium dichromate, acid dichromate, potassium permanganate, MnO₂, ruthenium tetroxide, phase transfer catalysts such as chromic acid or permanganate supported on a polymer, Cl₂-pyridine, H₂O₂-ammonium molybdate, NaO₂-CAN, NaOCl in HOAc, copper chromate, copper oxide, Raney nickel, palladium acetate, Meerwin-Pondorf-Verley reagent (aluminum *t*-butoxide with another ketone) and *N*-bromosuccinimide.


20 Then coupling of an organometallic carbon nucleophile such as a Grignard reagent, an organolithium, lithium dialkylcopper or R⁶-SiMe₃ in TAF with the ketone and an appropriate non-protic solvent at a suitable temperature, yields the 2'-alkylated sugar. The alkylated sugar optionally can be protected with a suitable protecting group, preferably with an acyl or silyl group, by methods well known to those skilled in the art, as taught by Greene et al., Protective Groups in Organic Synthesis, John Wiley and Sons, Second Edition, 1991.

25 The optionally protected sugar can then be coupled to the base by methods well known to those skilled in the art, as taught by Townsend, Chemistry of Nucleosides and Nucleotides, Plenum Press, 1994. For example, an acylated sugar can be coupled to a silylated base with a Lewis acid, such as tin tetrachloride, titanium tetrachloride, or trimethylsilyltriflate in an appropriate solvent at a suitable temperature. Alternatively, a halo-sugar can be coupled to a silylated base in the presence of trimethylsilyltriflate.

Subsequently, the nucleoside can be deprotected by methods well known to those skilled in the art, as by Greene et al., Protective Groups in Organic Synthesis, John Wiley and Sons, Second Edition, 1991.

In a particular embodiment, the 2'-C-branched ribonucleoside is desired, the synthesis of which is shown in Scheme 3. Alternatively, a deoxyribonucleoside is desired. To obtain these nucleosides, the formed ribonucleoside can optionally be protected by methods well known to those skilled in the art, as by Greene et al., Protective Groups in Organic Synthesis, John Wiley and Sons, Second Edition, 1991, and then the 2'-OH can be reduced with a suitable reducing agent. Optionally, the 2'-OH can be activated to facilitate reduction, such as, for example, by the Barton reduction.

Scheme 3



Modification of a pre-formed nucleoside


The key starting material for this process is an appropriately substituted nucleoside with a 2'-OH and 2'-H. The nucleoside can be purchased or can be prepared by any known means including standard coupling techniques. The nucleoside optionally
5 can be protected with suitable protecting groups, preferably with acyl or silyl groups, by methods well known to those skilled in the art, as described in Greene et al., Protective Groups in Organic Synthesis, John Wiley and Sons, Second Edition, 1991.

The appropriately protected nucleoside then can be oxidized with an appropriate oxidizing agent in a compatible solvent at a suitable temperature to yield the 2'-modified
10 sugar. Possible oxidizing agents include Jones' reagent (a mixture of chromic and sulfuric acids), Collins' reagent (dipyridine Cr(VI)oxide), Corey's reagent (pyridinium chlorochromate), pyridinium dichromate, acid dichromate, potassium permanganate, MnO₂, ruthenium tetroxide, phase transfer catalysts such as chromic acid or permanganate supported on a polymer, Cl₂-pyridine, H₂O₂-ammonium molybdate, NaO₂-
15 CAN, NaOCl in HOAc, copper chromate, copper oxide, Raney nickel, palladium acetate, Meerwin-Pondorf-Verley reagent (aluminum *t*-butoxide with another ketone) and *N*-bromosuccinimide.

Subsequently, the nucleoside can be deprotected by methods well known to those skilled in the art, as by Greene et al., Protective Groups in Organic Synthesis, John Wiley
20 and Sons, Second Edition, 1991.


In a particular embodiment, a 2'-C-branched ribonucleoside is desired, the synthesis of which is shown in Scheme 4. Alternatively, the deoxyribonucleoside may be desired. To obtain these nucleosides, the formed ribonucleoside optionally may be protected by methods well known to those skilled in the art, as by Greene et al.,
25 Protective Groups in Organic Synthesis, John Wiley and Sons, Second Edition, 1991, and then the 2'-OH can be reduced with a suitable reducing agent. Optionally, the 2'-OH can be activated to facilitate reduction such as, for example, by the Barton reduction.

Scheme 4



In another embodiment of the invention, the L-enantiomers are desired. These L-enantiomers corresponding to the compounds of the invention may be prepared following the same general methods given above, but beginning with the corresponding L-sugar or nucleoside L-enantiomer as the starting material.

3'-C-branched ribonucleosides of the following structures:



10 The key starting material for this process is an appropriately substituted sugar with a 3'-OH and a 3'-H, with an appropriate leaving group (LG) such as, for example, an acyl group or a halogen. The sugar can be purchased or can be prepared by any known means including standard epimerization, substitution, oxidation and/or reduction techniques. The substituted sugar then can be oxidized by an appropriate oxidizing agent in a compatible solvent at a suitable temperature to yield the 3'-modified sugar.

Then coupling of an organometallic carbon nucleophile such as a Grignard reagent, an organolithium, lithium dialkylcopper or R^6-SiMe_3 in TAF with the ketone and an appropriate non-protic solvent at a suitable temperature, yields the 3'-C-branched


sugar. The 3'-C-branched sugar optionally can be protected with a suitable protecting group, preferably with an acyl or silyl group, by methods well known to those skilled in the art, as taught by Greene et al., Protective Groups in Organic Synthesis, John Wiley and Sons, Second Edition, 1991.

5 The optionally protected sugar can then be coupled to the base by methods well known to those skilled in the art, as taught in Townsend, Chemistry of Nucleosides and Nucleotides, Plenum Press, 1994. For example, an acylated sugar can be coupled to a silylated base with a Lewis acid, such as tin tetrachloride, titanium tetrachloride, or trimethylsilyltriflate in an appropriate solvent at a suitable temperature. Alternatively, a
10 halo-sugar can be coupled to a silylated base in the presence of trimethylsilyltriflate.

Subsequently, the nucleoside can be deprotected by methods well known to those skilled in the art, as by Greene et al., Protective Groups in Organic Synthesis, John Wiley and Sons, Second Edition, 1991.

15 In a particular embodiment, the 3'-C-branched ribonucleoside is desired, the synthesis of which is shown in Scheme 5. Alternatively, a deoxyribonucleoside is desired. To obtain these nucleosides, the formed ribonucleoside can optionally be protected by methods well known to those skilled in the art, as by Greene et al., Protective Groups in Organic Synthesis, John Wiley and Sons, Second Edition, 1991, and then the 2'-OH can be reduced with a suitable reducing agent. Optionally, the 2'-
20 OH can be activated to facilitate reduction, such as, for example, by the Barton reduction.

Scheme 5



Modification of a pre-formed nucleoside.

5 The key starting material for this process is an appropriately substituted nucleoside with a 3'-OH and 3'-H. The nucleoside can be purchased or can be prepared by any known means including standard coupling techniques. The nucleoside can be optionally protected with suitable protecting groups, preferably with acyl or silyl groups, by methods well known to those skilled in the art, as taught by Greene et al., Protective Groups in Organic Synthesis, John Wiley and Sons, Second Edition, 1991.


10 The appropriately protected nucleoside can then be oxidized with the appropriate oxidizing agent in a compatible solvent at a suitable temperature to yield the 2'-modified

sugar. Possible oxidizing agents include Jones' reagent (a mixture of chromic and sulfuric acids), Collins' reagent (dipyridine Cr(VI)oxide), Corey's reagent (pyridinium chlorochromate), pyridinium dichromate, acid dichromate, potassium permanganate, MnO₂, ruthenium tetroxide, phase transfer catalysts such as chromic acid or permanganate supported on a polymer, Cl₂-pyridine, H₂O₂-ammonium molybdate, NaO₂-CAN, NaOCl in HOAc, copper chromate, copper oxide, Raney nickel, palladium acetate, Meerwin-Pondorf-Verley reagent (aluminum *t*-butoxide with another ketone) and *N*-bromosuccinimide.

Subsequently, the nucleoside can be deprotected by methods well known to those skilled in the art, as by Greene et al., Protective Groups in Organic Synthesis, John Wiley and Sons, Second Edition, 1991.

In a particular embodiment, the 3'-C-branched ribonucleoside is desired, the synthesis of which is shown in Scheme 6. Alternatively, a deoxyribonucleoside is desired. To obtain these nucleosides, the formed ribonucleoside can optionally be protected by methods well known to those skilled in the art, as by Greene et al.; Protective Groups in Organic Synthesis, John Wiley and Sons, Second Edition, 1991, and then the 2'-OH can be reduced with a suitable reducing agent. Optionally, the 2'-OH can be activated to facilitate reduction, such as, for example, by the Barton reduction.


Scheme 6



In another embodiment of the invention, the L-enantiomers are desired. These L-enantiomers corresponding to the compounds of the invention may be prepared following the same general methods given above, but beginning with the corresponding L-sugar or nucleoside L-enantiomer as the starting material.

General Synthesis of 4'-C-branched Nucleosides

4'-C-branched ribonucleosides of the following structures:



- 5 wherein R, R¹, R^{1'}, R², R^{2'}, R³, R^{3'}, X, X*, and Base are all as described above, can be prepared according to the following general methods.

Modification from the pentodialdo-furanose.

10 The key starting material for this process is an appropriately substituted pentodialdo-furanose. The pentodialdo-furanose can be purchased or can be prepared by any known means including standard epimerization, substitution and cyclization techniques.

15 In a preferred embodiment, the pentodialdo-furanose is prepared from the appropriately substituted hexose. The hexose can be purchased or can be prepared by any known means including standard epimerization (for eg., via alkaline treatment), substitution, and coupling techniques. The hexose can be in either the furanose form or cyclized by any means known in the art, such as methodology taught by Townsend in Chemistry of Nucleosides and Nucleotides, Plenum Press, 1994, preferably by selectively protecting the hexose, to give the appropriate hexafuranose.

20 The 4'-hydroxymethylene of the hexafuranose then can be oxidized with an appropriate oxidizing agent in a compatible solvent at a suitable temperature to yield the 4'-aldo-modified sugar. Possible oxidizing agents are Swern reagents, Jones' reagent (a mixture of chromic and sulfuric acids), Collins' reagent (dipyridine Cr(VI)oxide), Corey's reagent (pyridinium chlorochromate), pyridinium dichromate, acid dichromate, 25 potassium permanganate, MnO₂, ruthenium tetroxide, phase transfer catalysts such as

chromic acid or permanganate supported on a polymer, Cl₂-pyridine, H₂O₂-ammonium molybdate, NaO₂-CAN, NaOCl in HOAc, copper chromate, copper oxide, Raney nickel, palladium acetate, Meerwin-Pondorf-Verley reagent (aluminum *t*-butoxide with another ketone) and *N*-bromosuccinimide, although using H₃PO₄, DMSO and DCC in a mixture of benzene/pyridine at room temperature is preferred.

Then the pentodialdo-furanose optionally can be protected with a suitable protecting group, preferably with an acyl or silyl group, by methods well known to those skilled in the art, as taught by Greene et al., Protective Groups in Organic Synthesis, John Wiley and Sons, Second Edition, 1991. In the presence of a base, such as sodium hydroxide, the protected pentodialdo-furanose then can be coupled with a suitable electrophilic alkyl, halogeno-alkyl (such as CF₃), alkenyl or alkynyl (i.e., allyl), to obtain the 4'-alkylated sugar. Alternatively, the protected pentodialdo-furanose can be coupled with a corresponding carbonyl, such as formaldehyde, in the presence of a base like sodium hydroxide and with an appropriate polar solvent like dioxane, at a suitable temperature, and then reduced with an appropriate reducing agent to provide the 4'-alkylated sugar. In one embodiment, the reduction is carried out using PhOC(S)Cl and DMAP in acetonitrile at room temperature, followed by reflux treatment with ACCN and TMSS in toluene.

The optionally activated sugar can be coupled to the base by methods well known to those skilled in the art, as taught by Townsend in Chemistry of Nucleosides and Nucleotides, Plenum Press, 1994. For example, an acylated sugar can be coupled to a silylated base with a Lewis acid, such as tin tetrachloride, titanium tetrachloride, or trimethylsilyltriflate in an appropriate solvent at room temperature.

Subsequently, the nucleoside can be deprotected by methods well known to those skilled in the art, as by Greene et al., Protective Groups in Organic Synthesis, John Wiley and Sons, Second Edition, 1991.

In a particular embodiment, the 4'-C-branched ribonucleoside is desired. Alternatively, a deoxyribonucleoside is desired. To obtain these nucleosides, the formed ribonucleoside can optionally be protected by methods well known to those skilled in the art, as by Greene et al., Protective Groups in Organic Synthesis, John Wiley and Sons, Second Edition, 1991, and then the 2'-OH can be reduced with a suitable reducing agent.

Optionally, the 2'-OH can be activated to facilitate reduction, such as, for example, by the Barton reduction.

5 In another embodiment of the invention, the L-enantiomers are desired. These L-enantiomers corresponding to the compounds of the invention may be prepared following the same general methods given above, but beginning with the corresponding L-sugar or nucleoside L-enantiomer as the starting material.


Methods for Ribofuranosyl-2-azapurine Synthesis

10 *Preparation of 1'-C-methyl-ribofuranosyl-2-azapurine via 6-amino-9-(1-deoxy-beta-D-psicofuranosyl)purine.*

As an alternative method of preparation, the title compound can be prepared according to the published procedure of Farkas and Sorm (J. Farkas and F. Sorm, "Nucleic acid components and their analogues. XCIV. Synthesis of 6-amino-9-(1-deoxy-beta-D-psicofuranosyl)purine," *Collect. Czech. Chem. Commun.*, 1967, 32:2663-15 7; and J. Farkas, *Collect. Czech. Chem. Commun.*, 1966, 31:1535 (Scheme 7).

In a similar manner, but using the appropriate sugar and 2-azapurine base corresponding to the desired product compound, a variety of Formula (I) and/or Formula (II) compounds can be prepared.

Scheme 7




Alternative Methods for Ribofuranosyl-Purine Analogue Synthesis

Preparation of ribofuranosyl-purine analogues: 2-aza-3,7-dideazaadenosine derivative compounds.

Preparation of 2-aza-3,7-dideazaadenosine derivative compounds may be prepared according to the published synthesis of L.Townsend et al., *Bioorganic & Med. Chem. Letters*, 1991, 1(2):111-114, where the starting material, ethyl-3-cyanopyrrole-2-

carboxylate **4** was synthesized by Huisgen & Laschtuvka, according to the procedure provided in *Chemische Berichte*, 1960, 93:65-81, as shown in Scheme 8:


Scheme 8



Preparation of ribofuranosyl-purine analogues: 2-aza-3-deazaadenosine derivative compounds.


Preparation of 2-aza-3-deazaadenosine derivative compounds may be prepared according to the published synthesis of B. Otter et al., *J.Heterocyclic Chem.*, 1984, 481-89 shown in Scheme 9. The commercially available starting material used is the 4,5-dichloro-6-pyridazone 12.

Scheme 9



An alternative preparation of 2-aza-3-deazaadenosine derivative compounds that utilizes a chlorination step is that according to R.Panzica, *J. Chem. Soc. Perkin Trans I*, 1989, 1769-1774 and *J.Med.Chem.*, 1993, 4113-4120, shown in Scheme 10:


Scheme 10



5 *Preparation of purine analogues for nucleosides: optionally substituted 2,8-diaza-3,7-dideazaadenine derivative compounds.*

Preparation of certain 2,8-diaza-3,7-dideazaadenosine derivative compounds may be prepared according to the published synthesis by Oda et al. in *J.Heterocyclic Chem.*, 1984, 21:1241-55 and *Chem. Pharm. Bull.*, 1984, 32(11):4437-46, as shown in Scheme 11. The starting material is commercially available 4,5-dichloro-6-pyridazone 12.


Scheme 11



Preparation of purine analogues for nucleosides: 2,8-diaza-3-deazaadenosine derivative compounds.

Preparation of certain 2,8-diaza-3-deazaadenosine derivative compounds may be prepared according to the published synthesis by Panzica et al. in *J.Heterocyclic Chem.*, 1982, 285-88, *J. Med. Chem.*, 1993, 4113-20, and *Bioorg. & Med. Chem. Letters.*, 1996, 4(10):1725-31, as provided in Scheme 12. The key intermediate **27** was prepared via a 1,3-dipolar cycloaddition reaction between the 2,3,5-tri-*O*-benzoyl- β -D-ribofuranosyl azide **26** and methyl-hydroxy-2-butynoate **25**. A ribofuranosyl azide **26** synthesis was described by A. Stimac et al., *Carbohydrate Res.*, 1992, 232(2):359-65, using SnCl_4 catalyzed azidolysis of 1-*O*-Acetyl-2,3,5-tri-*O*-benzoyl- β -D-ribofuranose with Me_3SiN_3 in CH_2Cl_2 at room temperature.


Scheme 12



5 *Preparation of purine analogues for nucleosides: alternative preparation of 2,8-diaza-3-deazaadenine derivative compounds.*

2,8-diaza-3-deazaadenine derivative compounds may be prepared (see Scheme 13) according to the published synthesis by Chen et al. in *J.Heterocyclic Chem.*, 1982, 285-88; however, no condensation of this compound with ribofuranose is found.

Scheme 13



Preparation of ribofuranosyl-2-azapurines via use of protective groups.

5 As an alternative method of preparation, the compounds of the present invention can also be prepared by synthetic methods well known to those skilled in the art of nucleoside and nucleotide chemistry, such as taught by Townsend in Chemistry of Nucleosides and Nucleotides, Plenum Press, 1994.

10 A representative general synthetic method is provided in Scheme 14. The starting material is a 3,5-is-O-protected beta-D-alkyl ribofuranoside, but it will be understood that any 2', 3', or 5'-position may carry a protecting group to shield it from reacting. The 2'-C-OH then is oxidized with a suitable oxidizing agent in a compatible solvent at a suitable temperature to yield the 2'-keto-modified sugar. Possible oxidizing agents are Swern reagents, Jones' reagent (a mixture of chromic and sulfuric acids),
 15 Collins' reagent (dipyridine Cr(VI)oxide), Corey's reagent (pyridinium chlorochromate), pyridinium dichromate, acid dichromate, potassium permanganate, MnO_2 , ruthenium tetroxide, phase transfer catalysts such as chromic acid or permanganate supported on a polymer, Cl_2 -pyridine, H_2O_2 -ammonium molybdate, NaO_2 -CAN, NaOCl in HOAc, copper chromate, copper oxide, Raney nickel, palladium acetate, Meerwin-Pondorf-Verley reagent (aluminum *t*-butoxide with another ketone) and *N*-bromosuccinimide.

20


Next, addition of a Grignard reagent, such as, for example, an alkyl-, alkenyl- or alkynyl-magnesium halide like CH_3MgBr , $\text{CH}_3\text{CH}_2\text{MgBr}$, vinylMgBr , allylMgBr and

ethynylMgBr, or an alkyl-, alkenyl- or alkynyl-lithium, such as CH_3Li , in a suitable organic solvent, such as, for example, diethyl ether or THF, across the double bond of the 2'-carbonyl group provides a tertiary alcohol at this position. The addition of a hydrogen halide in a suitable solvent, such as, for example, H₂ in HOAc, in the subsequent step provides a leaving group (LG) such as, for example, a chloro, bromo or iodo, at the C-1 anomeric carbon of the sugar ring that later generates a nucleosidic linkage. Other suitable LGs include C-1 sulfonates such as, for example, methanesulfonate, trifluoromethanesulfonate and/or *p*-toluenesulfonate.

The introduction in the next step of a metal salt (Li, Na or K) of an appropriately substituted 2-azapurine in a suitable organic solvent such as, for example, THF, acetonitrile or DMF, results in the formation of the desired nucleosidic linkage and addition of the desired 2-azapurine base. This displacement reaction may be catalyzed by a phase transfer catalyst like TDA-1 or triethylbenzylammonium chloride. The introduction of a "Z" substituent on any of base formulae (i)-(vi) optionally may be performed subsequent to the initial addition of protecting groups. For example, the introduction of an amino group for "Z" is accomplished by the addition of an appropriate amine in an appropriate solvent to the 2'-C-halo intermediate just prior to the last step of removal of the protecting groups. Appropriate amines include alcoholic or liquid ammonia to generate a primary amine (-NH₂), an alkylamine to generate a secondary amine (-NHR), or a dialkylamine to generate a tertiary amine (-NRR').

Finally, the nucleoside can be deprotected by methods well known to those skilled in the art, as by Greene et al., Protective Groups in Organic Synthesis, John Wiley and Sons, Second Edition, 1991. It is to be noted that this reaction scheme can be used for joining any of the purine nucleoside analogue bases provided for in Schemes 8-13 with a ribofuranosyl moiety.

Scheme 14



The present invention is described by way of illustration in the following examples. It will be understood by one of ordinary skill in the art that these examples are in no way limiting and that variations of detail can be made without departing from the spirit and scope of the present invention.

EXAMPLES

The test compounds were dissolved in DMSO at an initial concentration of 200 μ M and then were serially diluted in culture medium.

Unless otherwise stated, bay hamster kidney (HK-21) (ATCC CCL-10) and bos Taurus (T) (ATCC CRL 1390) cells were grown at 37°C in a humidified CO₂ (5%) atmosphere. HK-21 cells were passaged in Eagle MEM additioned of 2 mM L-glutamine, 10% fetal ovine serum (FS, Gibco) and Earle's SS adjusted to contain 1.5 g/L sodium bicarbonate and 0.1 mM non-essential amino acids. T cells were passaged in Dulbecco's modified Eagle's medium with 4 mM L-glutamine and 10% horse serum (HS, Gibco), adjusted to contain 1.5 g/L sodium bicarbonate, 4.5 g/L glucose and 1.0 mM sodium pyruvate. The vaccine strain 17D (YFV-17D) (Stamaril®, Pasteur Merieux) and Bovine Viral Diarrhea virus (BVDV) (ATCC VR-534) were used to infect HK and T cells, respectively, in 75 cm² bottles. After a 3 day incubation period at 37°C, extensive cytopathic effect was observed. Cultures were freeze-thawed three times, cell debris were removed by centrifugation and the supernatant was aliquoted and stored at -70°C. YFV-17D and VDV were titrated in HK-21 and T cells, respectively, that were grown to confluency in 24-well plates.

The following examples are derived by selection of an appropriate, optionally substituted sugar or cyclopentane ring coupled with an optionally substituted 2-azapurine base, and prepared according to the following synthetic schemes:

Example 1: Synthesis of optionally substituted 1'-C-branched-ribofuranosyl, -sulfonyl, -thiophenyl or cyclopentanyl-2-azapurines;

Example 2: Synthesis of optionally substituted 2'-C-branched -ribofuranosyl, -sulfonyl, -thiophenyl or cyclopentanyl-2-azapurines;

Example 3: Synthesis of optionally substituted 3'-C-branched-ribofuranosyl, -sulfonyl, -thiophenyl or cyclopentanyl-2-azapurines;


Example 4: Synthesis of optionally substituted 4'-C-branched -ribofuranosyl, -sulfonyl, -thiophenyl or cyclopentanyl-2-azapurines;

Examples 5 – 13: Synthesis of specific compounds of the present invention; and

Examples 14 – 18: Biologic test results of representative examples of compounds of the present invention.

Example 1. 1'-C-branched ribofuranosyl, -sulfonyl or cyclopentanyl-2-azapurine, optionally substituted.


The title compound is prepared according to Schemes 1, 2, or 7. In a similar manner but using the appropriate sugar or cyclopentane ring and optionally substituted 2-azapurine base, the following nucleosides of Formulae (I) or (II) may be prepared:




wherein: base may be any of the Formulae (A)-(G) as described herein where R in each instance may exist in mono-, di- or triphosphate form.

Alternatively, the Dimroth rearrangement may be used for making 2-azapurines from the corresponding purine base. In this reaction, an N-alkylated or N-arylated imino heterocycle undergoes rearrangement to its corresponding alkylamino or arylamino heterocycle.

Example 1a: 1'-C-hydroxymethyl-2-azaadenosine



Example 2a: 2'-C-methyl-2-azaadenosine



(Synthesis according to the procedure of J. A. Montgomery, *Nucleic Acid Chemistry*, 1978, Part II, 681-685 starting with 2'-C-methyladenosine.)

5

Step 1: H_2O_2 , AcOH, 80%; Step 2: BnBr, DMAc, Step 3: NaOH, H_2O , EtOH, 30%, Step 4: NH_3/MeOH , 80°C , 2 days, 60%; Step 5: $\text{H}_2/\text{Pd/C}$, 3 atm, MeOH, 30% Step 6: NaNO_2 , AcOH, H_2O , 50%.


10 4-Amino-7-(2-C-methyl- β -D-ribofuranosyl)imidazo[4,5-d]-v-triazine (2'-C-methyl-2-azaadenosine): ^1H NMR ($\text{DMSO}-d_6$) δ 8.82 (s, 1H, H8), 7.97 (br, 2H, NH_2), 6.12 (s, 1H, H1'), 5.22-5.51 (m, 3H, 3OH), 3.70-4.17 (m, 4H, H3', H4', 2H5'), 0.80 (s, 3H, CH_3). ^{13}C NMR ($\text{DMSO}-d_6$) δ 153, 146, 142, 116, 92, 83, 79, 72, 60, 20. m/z (FAB >0) 565 $(2\text{M}+\text{H})^+$, 283 $(\text{M}+\text{H})^+$, (FAB <0) 563 $(2\text{M}-\text{H})^-$.

15

Alternatively, 2-azaadenosine shown as the final product in Example 2.a. may be prepared starting with adenosine, according to the procedure of J. A. Montgomery, *Nucleic Acid Chemistry*, 1978, Part II, 681-685 starting with 2'-C-methyladenosine, or via 2-azainosine in a synthetic procedure taught by R. P. Panzica, *Journal of Heterocyclic Chemistry*, 1972, 9:623- 628 starting with AICA riboside.

20


Example 2b: 2'-C-methyl-pyrrolo-4-amino-1,2,3-triazine



Step 1 : NCS, DMF; Step 2: mCPBA, AcOH; Step 3: a) BnBr, DMAc; b) NaOH, H₂O, EtOH, Step 4: NH₃/MeOH, 80°C, Step 5: H₂/Pd/C, MeOH; Step 6: .NaNO₂, AcOH, H₂O.

Example 3. 3'-C-branched ribofuranosyl, -sulfonyl or cyclopentanyl-2-azapurine, optionally substituted.

The title compound is prepared according to Schemes 5, 6, or through protection of appropriately selected substituent groups in Scheme 8. In a similar manner ut using the appropriate sugar or cyclopentane ring and optionally substituted 2-azapurine base, the following nucleosides of Formulae (I) and (II) may be prepared:




wherein: base may be any of the Formulae (A)-(G) as described herein where R in each instance may exist in mono-, di- or triphosphate form.

Alternatively, the Dimroth rearrangement may be used for making 2-azapurines from the corresponding purine base. In this reaction, an N-alkylated or N-arylated imino heterocycle undergoes rearrangement to its corresponding alkylamino or arylamino heterocycle.

Example 4. 4'-C-branched ribofuranosyl, -sulfonyl or cyclopentany-2-azapurine, optionally substituted.


The title compound is prepared according to modification from the corresponding pentodialdo-furanose. In a similar manner but using the appropriate sugar or cyclopentane ring and optionally substituted 2-azapurine base, the following nucleosides of Formulae (I) or (II) may be prepared:



wherein: base may be any of the Formulae (A)-(G) as described herein where R in each instance may exist in mono-, di- or triphosphate form.

Alternatively, the Dimroth rearrangement may be used for making 2-azapurines from the corresponding purine base. In this reaction, an N-alkylated or N-arylated imino heterocycle undergoes rearrangement to its corresponding alkylamino or arylamino heterocycle.

Example 5: Synthesis of 4-amino-1-(β -D-ribofuranosyl)imidazo[4,5-d] pyridazine



Step A : 1-(2,3,5-tri-*O*-Benzoyl- β -D-ribofuranosyl)-5-benzyloxymethylimidazo[4,5-*d*]pyridazin-4-one:

5 The 5-benzyloxymethylimidazo[4,5-*d*]pyridazine (500 mg, 1.95 mmol) [for preparation see Journal of Heterocyclic Chemistry, 1984, Vol 21, 481] was heated at reflux in hexamethyldisilazane (6 mL) for 1 hour. The mixture was evaporated to dryness to give a slight yellow syrup which was dissolved in dry 1,2-dichloroethane (20 mL). The 1-*O*-acetyl -2,3,5-tri-*O*-benzoyl- β -D-ribofuranose (1.04 g, 2.06 mmol) and stannic chloride (0.4 mL, 3.44 mmol) were added at 20°C and the mixture was stirred for 3 hours. The reaction mixture was poured into an aqueous solution of sodium hydrogenocarbonate, filtrated through a pad of celite and washed by dichloromethane. The organic layer was evaporated to dryness to give a yellow foam. The crude product was purified on silica gel using n-hexane/ethyl acetate (3/2) as eluant to give the title compound (703 mg) as a white powder.

¹H NMR (DMSO-*d*₆) δ ppm: 4.39 (s, 2H, CH₂), 4.60 (m, 2H), 4.73 (m, 1H), 5.34 (dd, 2H, CH₂), 5.77-5.88 (m, 2H, H2' and H3'), 6.56 (m, 1H, H1'), 6.98-7.10 (m, 5H), 7.23-7.32 (m, 6H), 7.41-7.51 (m, 3H), 7.68-7.73 (m, 2H), 7.74-7.8 (m, 4H), 8.51 (s, 1H), 8.52 (s, 1H).

Step B : 1-(2,3,5-tri-*O*-Benzoyl- β -D-ribofuranosyl)imidazo[4,5-*d*]pyridazin-4-one:

To a solution containing the compound from Step B (500 mg, 0.7 mmol), in dry dichloromethane (25 mL) was added a pre-cooled (-78°C) solution of boron trichloride

1M (5 mL) at -78°C and stirred for 2 hours at -78°C. A mixture of methanol/dichloromethane (1/1) was added to the mixture at -78°C and then at 20°C. The reaction mixture was evaporated to dryness to give a yellow powder. The crude product was purified on silica gel using n-hexane/ethyl acetate (3/2) as eluant to give the title compound (400 mg) as a yellow powder.

¹H NMR (DMSO-d₆) δ ppm: 4.77-4.98 (m, 3H, H4', 2H5'), 5.95-6.12 (m, 2H, H2' and H3'), 6.65 (m, 1H, H1'), 7.39-7.76 (m, 9H), 7.84-8.06 (m, 6H), 8.64-5.79 (m, 2H, H3 and H8), 12.84 (br, 1H, NH).

Mass spectrum: *m/z* (FAB>0) 581 (M+H)⁺, (FAB<0) 579 (M-H)⁻

Step C : 4-chloro-1-(2,3,5-tri-*O*-Benzoyl-β-D-ribofuranosyl)imidazo[4,5-*d*]pyridazine:

A solution containing the compound from Step B (1.32 g, 2.27 mmol), the *N,N*-diethylaniline (365 μL), tetrabutylammonium chloride (1.2 g), freshly distilled phosphorus chloride (1.3 μL) and anhydrous acetonitrile (17 mL) was stirred at 90°C for 1 hour. The reaction mixture was poured over cracked ice/water. The aqueous layer was extracted with dichloromethane (3x60 mL). The organic layer was washed with sodium hydrogenocarbonate 5%, water and was evaporated to dryness. The crude product was purified on silica gel using n-hexane/ethyl acetate (3/1) as eluant to give the title compound (404 mg) as a yellow powder.


¹H NMR (DMSO-d₆) δ ppm: 4.82-6.87 (m, 2H), 4.9-6.95 (m, 1H), 6.0-6.08 (m, 1H), 6.12-6.19 (m, 1H), 6.90 (d, 1H, *J* = 5.2 Hz, H1'), 7.47-7.73 (m, 9H), 7.88-8.12 (m, 6H), 9.10 (s, 1H, H8), 9.90 (s, 1H, H3)

Step D : 4-amino-1-(β-D-ribofuranosyl)imidazo[4,5-*d*]pyridazine:

The compound from Step C (420 mg, 0.7 mmol) was added to a solution of ammonia in methanol and stirred in a steel bomb at 150°C for 6 hours. The reaction mixture was evaporated to dryness to afford a brown oil which was purified on silica gel reverse-phase (C18) using water as eluant to give the title compound (50 mg) as a yellow powder.

^1H NMR (DMSO- d_6) δ ppm: 3.58-4.48 (m, 5H, H2', H3', H4', 2H5'), 5.14-5.68 (m, 3H, 3xOH), 5.90 (s, 1H, H1'), 6.61 (br, 2H, NH₂), 8.59 (s, 1H, H8), 9.12 (s, 1H, H3)

Example 6. Synthesis of 1-(β -D-ribofuranosyl)imidazo[4,5-*d*]pyridazin-4-one



5

1-(2,3,5-tri-*O*-Benzoyl- β -D-ribofuranosyl)imidazo[4,5-*d*]pyridazin-4-one (555 mg, 0.9 mmol) was added to a solution of sodium methylate (205 mg) in methanol (25 mL) and stirred at 20°C for 2 hours. The reaction mixture was evaporated to dryness. The residue was dissolved in water and washed with ethyl acetate. The aqueous layer was concentrated under pressure. The crude product was purified on silica gel reverse-phase (C18) using water as eluant to give the title compound (220 mg) as a white powder.

10


^1H NMR (DMSO- d_6) δ ppm: 3.59-3.62 (m, 2H), 4.02 (m, 1H), 4.11 (m, 1H), 4.22 (m, 1H), 5.16-5.72 (m, 3H, 3xOH), 5.91 (s, 1H, H1'), 8.52 (s, 1H, H8), 8.68 (s, 1H, H3), 12.75 (br, 1H, NH).

15

Mass spectrum: m/z (FAB>0) 537 (2M+H)⁺, 269 (M+H)⁺, (FAB<0) 535 (2M+H)⁺, 267 (M-H)⁻

Example 7. Synthesis of 4-amino-1-(2-*C*-methyl- β -D-ribofuranosyl)imidazo[4,5-*d*]pyridazine

20



Step A : 1-(2-C-methyl-2,3,5-tri-O-Benzoyl- β -D-ribofuranosyl)imidazo[4,5-*d*]pyridazin-4-one

To a suspension of imidazo[4,5-*d*]pyridazine (3.48 g, 25.5 mmol) [for preparation see Journal of Heterocyclic Chemistry, 1969, Vol 6, 93] in dry acetonitrile (35 mL) was added 1,2,3,5-tetra-O-benzoyl-2-C-methyl- β -D-ribofuranose (14.48 g, 25.0 mmol) at 20°C and stirred for 15 mn. DBU (11.5 mL, 76.3 mmol) was added at 0°C and the solution was stirred for 15 mn at 0°C. TMSOTf (24.7 mL, 127.8 mmol) was added at 0°C and the mixture was heated at 80°C for 20 hours. The reaction mixture was poured into an aqueous solution of sodium hydrogenocarbonate and extracted by ethyl acetate. The organic layer was evaporated to dryness to give a yellow powder. The crude product was purified on silica gel using dichloromethane/methanol (99.3/0.7) as eluant to give a slight yellow powder which was crystallized from isopropanol to give the title compound (2.45 g) as a white powder.

¹H NMR (DMSO-*d*₆) δ ppm: 1.48 (s, 3H, CH₃), 4.75-4.96 (m, 3H, H4', 2H5'), 5.81 (d, 1H, J= 5.5 Hz, H3'), 6.99 (s, 1H, H1'), 7.39-7.72 (m, 9H), 7.92-8.08 (m, 6H), 8.64 (s, 1H, H8), 8.71 (s, 1H, H3), 12.89 (br, 1H, NH).

Mass spectrum: *m/z* (FAB>0) 1189 (2M+H)⁺, 585 (M+H)⁺, (FAB<0) 593 (M-H)⁻

Step B: 4-chloro-1-(2-C-methyl-2,3,5-tri-O-Benzoyl- β -D-ribofuranosyl)imidazo[4,5-*d*]pyridazine:

A solution containing the compound from Step A (300 mg, 0.50 mmol), the N,N-diethylaniline (1.2 mL) and freshly distilled phosphorus chloride (24 mL) was stirred at reflux for 1 hour. The reaction mixture was evaporated to dryness. Dichloromethane was added to the residue and the organic layer poured over cracked ice/water. The aqueous layer was extracted with dichloromethane. The organic layer was washed with sodium hydrogenocarbonate 5%, water and was evaporated to dryness. The crude product was purified on silica gel using diethyl ether/petrol ether (1/1) as eluant to give the title compound (295 mg) as a white powder.

¹H NMR (DMSO-d₆) δ ppm: 1.5 (s, 3H, CH₃), 4.8-5.0 (m, 3H, H4', 2H5'), 5.85 (d, 1H, J= 5.5 Hz, H3'), 7.15 (s, 1H, H1'), 7.38-8.08 (m, 15H), 9.15 (s, 1H, H8), 9.90 (s, 1H, H3)

5 Step C: 4-amino-1-(2-C-methyl-β-D-ribofuranosyl)imidazo[4,5-d]pyridazine:


The compound from Step B (590 mg, 0.96 mmol) was added to a solution of ammonia in methanol and stirred in a steel bomb at 150°C for 6 hours. The reaction mixture was evaporated to dryness to remove methanol. The crude product was purified on silica gel reverse-phase (C18) using water as eluant to give the title compound (35
10 mg) as a white powder.

¹H NMR (DMSO-d₆) δ ppm: 0.70 (s, 3H, CH₃), 3.64-3.98 (m, 4H, H3', H4', 2H5'), 5.23-5.44 (m, 3H, 3OH), 5.98 (s, 1H, H1'), 6.63 (br, 2H, NH₂), 8.68 (s, 1H, H8), 9.05 (s, 1H, H3)

¹³C NMR (DMSO-d₆) δ ppm: 155, 143, 132, 131, 129, 93, 83, 79, 72, 20.

15 Mass spectrum: *m/z* (FAB>0) 282 (M+H)⁺, (FAB<0) 280 (M-H)⁻

Example 8: Synthesis of the 4-substituted-1-(2-C-methyl- β -D-ribofuranosyl)imidazo[4,5-*d*]pyridazine



W	Step A	products
OH	NaOMe, MeOH 100°C, 24 h	
Cl	NH ₃ /MeOH 20°C, 48 h	

5 **1-(2-C-methyl- β -D-ribofuranosyl)imidazo[4,5-*d*]pyridazin-4-one:**

¹H NMR (DMSO-*d*₆) δ ppm: 1.17 (s, 3H, CH₃), 3.44-3.59 (m, 1H), 3.68-3.78 (m, 1H), 3.86-3.94 (m, 1H), 4.11-4.21 (m, 1H), 4.8-5.4 (m, 3H, 3OH), 6.05 (s, 1H, H1'), 8.35 (s, 1H, H8), 8.37 (s, 1H, H3), 12.67 (br, 1H, NH).

Mass spectrum: m/z (FAB>0) 283 (2M+H)⁺, 281 (M+H)⁺

10


4-chloro-1-(2-C-methyl- β -D-ribofuranosyl)imidazo[4,5-*d*]pyridazine

¹H NMR (DMSO-*d*₆) δ ppm: 0.72 (s, 3H, CH₃), 3.69-4.06 (m, 4H, H3', H4', 2H5'), 5.34-5.51 (m, 3H, 3OH), 6.19 (s, 1H, H1'), 9.18 (s, 1H, H8), 9.87 (s, 1H, H3)

Mass spectrum: m/z (FAB>0) 301 (M+H)⁺, (FAB<0) 299 (M-H)⁻

15

Example 9: Synthesis of the 4,7-diamino-imidazo[4,5-*d*]pyridazine nucleosides derivatives







Step A:

5 To a suspension of 4,5-dicyanoimidazole (1 eq.) [for preparation see Journal of Organic Chemistry, 1976, Vol 41, 713] in dry DMF (0.2 M) was added the protected β -D-ribofuranose derivatives (1 eq.) at 20°C. DBU (3 eq.) was added at 0°C and the solution was stirred for 20 mn at 0°C. TMSOTf (4 eq.) was added at 0°C and the mixture was heated at 60°C for 1 hour. The reaction mixture was poured into an aqueous solution of sodium hydrogenocarbonate and extracted by dichloromethane. The organic layer was evaporated to dryness to give a yellow powder. The crude product was purified on silica gel using diethyl ether/petrol ether as eluant to give the title compound (see the following table 1).

15 **Step B:**

 The compound from Step A (1 eq.) was stirred with hydrazine monohydrate (20 eq.) and acetic acid (1.4 eq.) at 75°C for several hours (see the following table 1). The reaction mixture was poured into water. The aqueous layer was washed by dichloromethane and evaporated under pressure. The residue was purified on reverse-phase column to give the title compound (see the following table 1).

Table 1:

R	products from Step A	yield (Step A)	experiments (Step B)	products from Step B	yield (Step A)
H		63 %	75°C for 1.5h.		58 % (white powder)
CH ₃		62 %	75°C for 20h.		15 % (white powder)

4,7-diamino-1-β-D-ribofuranosylimidazo[4,5-d]pyridazine:

- 5 ¹H NMR (DMSO-d₆) δ ppm: 3.58–4.32 (m, 5H, H2', H3', H4', 2H5'), 5.10–5.90 (br, 7H, 2NH₂, 3OH), 6.11 (s, 1H, H1'), 8.50 (s, 1H, H8)

¹³C NMR (DMSO-d₆) δ ppm: 151, 144, 142, 132, 122, 89, 86, 75, 70, 61.

Mass spectrum: *m/z* (FAB>0) 283 (M+H)⁺, (FAB<0) 281 (M-H)⁻


- 10 **4,7-diamino-1-(2-C-methyl-β-D-ribofuranosyl)imidazo[4,5-d]pyridazine:**

¹H NMR (DMSO-d₆) δ ppm: 0.75 (s, 3H, CH₃), 3.67–3.76 (m, 1H), 3.84–3.94 (m, 3H), 5.32 (m, 3H, 3OH), 5.43 (br, 1H, NH₂), 5.71 (br, 1H, NH₂), 6.21 (s, 1H, H1'), 8.78 (s, 1H, H8)

¹³C NMR (DMSO-d₆) δ ppm: 151, 144, 142, 132, 123, 92, 83, 78, 71, 59, 20.

- 15 Mass spectrum: *m/z* (FAB>0) 593 (2M+H)⁺, 297 (M+H)⁺, (FAB<0) 295 (M-H)⁻

Example 10: Synthesis of 4,7-disubstituted-imidazo[4,5-*d*]pyridazine nucleosides



Step A : Typical procedure for the preparation of the protected 4,7-dichloroimidazo[4,5-*d*] pyridazine nucleosides:

The 4,7-dichloroimidazo[4,5-*d*]pyridazine [for preparation see Journal of Heterocyclic Chemistry, 1968, Vol 5, 13] (1 eq.) was heated at reflux in hexamethyldisilazane for 12 hours. The mixture was evaporated to dryness to give a solid which was dissolved in 1,2-dichloroethane. The protected β -D-ribofuranose derivatives (1.1 eq.) and stannic chloride (1.4 eq.) were added at 20C and the solution was stirred for 3 hours. The reaction mixture was poured into an aqueous solution of sodium hydrogenocarbonate, filtrated through a pad of celite and washed by dichloromethane. The organic layer was evaporated to dryness. The crude product was purified on silica gel using dichloromethane /acetone (40/1) as eluant to give the title compound (see the following table 2).

Step B: Typical procedure for the preparation of the 4,7-dichloroimidazo[4,5-*d*]pyridazine nucleosides:

5 The compound from Step A (1 eq.) was stirred with sodium methoxide (0.1 eq.) in methanol for several hours. The reaction mixture was evaporated under pressure. Water was added to the residue. The aqueous layer was washed by ethyl acetate and was evaporated under pressure. The residue was purified on reverse-phase column to give the title compound (see the following table 2).

10 Step C: Typical procedure for the preparation of the imidazo[4,5-*d*]pyridazine nucleosides:

15 A mixture of the compound from Step A (1 eq.), palladium on charcoal (10%), sodium acetate (4.2 eq.) in acetyl acetate was stirred under hydrogen until the compound from Step A was consumed. The reaction mixture was evaporated under pressure and was purified on silica gel to give the title protected compound which was stirred with sodium methoxide (3.3 eq.) in methanol. The reaction mixture was evaporated under pressure. Water was added to the residue. The aqueous layer was washed by ethyl acetate and was evaporated under pressure. The residue was purified on reverse-phase column to give the title compound (see the following table 2).

20 Step D: Typical procedure for the preparation of the chloro-methoxy-imidazo[4,5-*d*]pyridazine nucleosides:

25 The compound from Step A (1 eq.) was stirred with sodium methoxide (3.3 eq.) in methanol 0.3M at 20°C for several hours. The reaction mixture was evaporated under pressure. Water was added to the residue. The aqueous layer was washed by ethyl acetate and was evaporated under pressure. The residue was purified on reverse-phase column to give a compound whose regioselectivity was not given (see the following table 2).

Step E: Typical procedure for the preparation of the methoxy-imidazo[4,5-*d*]pyridazine nucleosides:

5 A mixture of the compound from Step D (1 eq.), palladium on charcoal (10%), sodium acetate (4.2 eq.) in water/ethanol (1/1) was stirred under hydrogen until the compound from Step A was consumed. The reaction mixture was evaporated under pressure and was purified on reverse-phase column to give the title compound whose regioselectivity was not given (see the following table 2).

10 Step F: Typical procedure for the preparation of the protected 4,7-diazidoimidazo[4,5-*d*]pyridazine nucleosides:

15 The compound from Step A (1 eq.) was treated at 50°C with sodium azide (1.5 eq.) in DMF. Water was added to the mixture. The aqueous layer was extracted by ethyl acetate. The organic layer was evaporated under pressure. The crude product was purified on silica gel using diethyl ether/petrol ether (7/3) as eluant to give the title compound (see the following table 2).

Step G: Typical procedure for the preparation of the azido-methoxy-imidazo[4,5-*d*]pyridazine nucleosides:






20 The compound from Step F (1 eq.) was stirred at 50°C with sodium methoxide (1 eq.) in methanol. The reaction mixture was evaporated under pressure. Water was added to the residue. The aqueous layer was washed by ethyl acetate and was evaporated under pressure. The residue was purified on reverse-phase column using water/acetonitrile as eluant to give the title compound whose regioselectivity was not given (see the following table 2).

25 Step H: Typical procedure for the preparation of the amino-azido-imidazo[4,5-*d*]pyridazine nucleosides:

A mixture of the compound from Step F (1 eq.), palladium on charcoal (10%), sodium acetate (4.2 eq.) in ethyl acetate was stirred under hydrogen until the compound

from Step F was consumed. The reaction mixture was filtrated over celite and was evaporated under pressure The crude product was purified on silica gel to give the title protected compound whose regioselectivity was not given (see the following table 1). This compound was stirred with sodium methoxide (3 eq.) in methanol. The reaction mixture was evaporated under pressure. Water was added to the residue. The aqueous layer was washed by ethyl acetate and was evaporated under pressure. The residue was purified on reverse-phase column to give the title compound whose regioselectivity was not given (see the following table 2).

Table 2:

experiments	products	yield
Step A		60 % (white powder)
Step A		34 % (white powder)
Step B		(white powder)
Step C		71 % (white powder)
Step D		55 % (white powder)

experiments	products	yield
Step E		55 % (white powder)
Step F		50 % (yellow powder)
Step F		50 % (yellow powder)
Step G		38 % (beige powder)
Step H		98 % (white powder)
Step H		49 % (white powder)

4,7-dichloro-1-(2,3,5-tri-O-Benzoyl- β -D-ribofuranosyl)imidazo[4,5-d]pyridazine:

^1H NMR (DMSO- d_6) δ ppm: 4.8-5.0 (m, 3H, H4', 2H5'), 6.05 (s, 1H, H3'), 6.25 (s, 1H, H2'), 7.1 (d, 1H, J = 4 Hz, H1'), 7.4-8.0 (m, 15H), 9.25 (s, 1H, H8).

Mass spectrum: m/z (FAB>0) 633 (M+H) $^+$

4,7-dichloro-1-(2-C-methyl-2,3,5-tri-O-Benzoyl- β -D-ribofuranosyl)imidazo[4,5-
d]pyridazine:

^1H NMR (DMSO- d_6) δ ppm: 1.65 (s, 3H, CH₃), 4.9-5.0 (m, 3H, H4', 2H5'), 5.8 (s, 1H, H3'), 7.35-8.05 (m, 16H including H1'), 9.3 (s, 1H, H8).

5

4,7-dichloro-1-(2-C-methyl- β -D-ribofuranosyl)imidazo[4,5-d]pyridazine:

^1H NMR (DMSO- d_6) δ ppm: 0.84 (s, 3H, CH₃), 3.77 (m, 1H), 3.88-4.04 (m, 3H), 5.30-5.60 (m, 3H, OH), 6.5 (s, 1H, H1'), 9.44 (s, 1H, H8).

10 1-(2-C-methyl- β -D-ribofuranosyl)imidazo[4,5-d]pyridazine:

^1H NMR (DMSO- d_6) δ ppm: 0.80 (s, 3H, CH₃), 3.75 (m, 1H), 3.80-4.00 (m, 3H), 5.40 (br, 3H, OH), 6.2 (s, 1H, H1'), 9.0 (s, 1H), 9.65 (s, 1H), 9.85 (s, 1H).

15 7-chloro-4-methoxy-1- β -D-ribofuranosylimidazo[4,5-d]pyridazine or 4-chloro-7-methoxy-1- β -D-ribofuranosylimidazo[4,5-d]pyridazine:

^1H NMR (DMSO- d_6) δ ppm: 3.6-4.5 (m, 8H, 2H5', H4', H3', H2', OCH₃), 5.40 (m, 3H, OH), 6.2 (s, 1H, H1'), 9.0 (s, 1H, H8).

675 (2M+H)⁺, Mass spectrum: m/z (FAB>0) 317 (M+H)⁺, (FAB<0) 315 (M-H)⁻

20 4-methoxy-1- β -D-ribofuranosylimidazo[4,5-d]pyridazine or 7-methoxy-1- β -D-ribofuranosyl imidazo[4,5-d]pyridazine:

^1H NMR (DMSO- d_6) δ ppm: 3.54-3.79 (m, 2H), 3.95 (m, 1H), 4.15 (m, 1H, H3'), 4.2 (s, 3H, OCH₃), 4.4 (m, 1H, H2'), 5.05-5.70 (m, 3H, OH), 6.2 (d, 1H, J = 4.8 Hz, H1'), 8.95 (s, 1H), 9.3 (s, 1H).

25 ^{13}C NMR (DMSO- d_6) δ ppm: 154, 145, 143, 142, 121, 90, 85, 75, 70, 61, 55.

Mass spectrum: m/z (FAB>0) 283 (M+H)⁺, (FAB<0) 281 (M-H)⁻

4,7-diazido-1-(2,3,5-tri-*O*-Benzoyl- β -D-ribofuranosyl)imidazo[4,5-*d*]pyridazine:

^1H NMR (DMSO- d_6) δ ppm: 4.81-5.1 (m, 3H, H4', 2H5'), 6.24-6.49 (m, 2H, H2', H3'), 7.2 (d, 1H, $J = 5$ Hz, H1'), 7.4-8.0 (m, 15H), 9.12 (s, 1H, H8).

5 Mass spectrum: m/z (FAB>0) 647 ($\text{M}+\text{H}$) $^+$

4,7-diazido-1-(2-*C*-methyl-2,3,5-tri-*O*-Benzoyl- β -D-ribofuranosyl)imidazo[4,5-*d*]pyridazine:

10 ^1H NMR (DMSO- d_6) δ ppm: 1.6 (s, 3H, CH_3), 4.96 (m, 3H, H4', 2H5'), 6.02 (m, 1H, H3'), 7.24 (s, 1H, H1'), 7.40-7.52 (m, 6H), 7.60-7.71 (m, 3H), 7.93-8.1 (m, 6H), 9.10 (s, 1H, H8).

Mass spectrum: m/z (FAB>0) 661 ($\text{M}+\text{H}$) $^+$

15 4-azido-7-methoxy-(2-*C*-methyl- β -D-ribofuranosyl)imidazo[4,5-*d*]pyridazine or 7-azido-4-methoxy-1-(2-*C*-methyl- β -D-ribofuranosyl)imidazo[4,5-*d*]pyridazine:

^1H NMR (DMSO- d_6) δ ppm: 0.75 (s, 3H, CH_3), 3.70-3.98 (m, 2H), 4.05 (m, 2H), 4.2 (s, 3H, OCH_3), 5.32-5.61 (br, 3H, OH), 6.36 (d, 1H, $J = 5.7$ Hz, H1'), 9.18 (s, 1H, H8),

^{13}C NMR (DMSO- d_6) δ ppm: 156, 143, 136, 129, 123, 94, 83, 79, 71, 59, 57, 20.

20 Mass spectrum: m/z (FAB>0) 675 ($2\text{M}+\text{H}$) $^+$, 338 ($\text{M}+\text{H}$) $^+$, (FAB<0) 673 ($2\text{M}-\text{H}$) $^-$, 336 ($\text{M}-\text{H}$)

4-amino-7-azido-1-(2-C-methyl-2,3,5-tri-O-Benzoyl- β -D-ribofuranosyl)imidazo[4,5-*d*]pyridazine or 7-amino-4-azido-1-(2-C-methyl-2,3,5-tri-O-Benzoyl- β -D-ribofuranosyl)imidazo[4,5-*d*]pyridazine:

¹H NMR (DMSO-*d*₆) δ ppm: 01.64 (s, 3H, CH₃), 4.95 (m, 3H), 6.06 (m, 1H, H3'), 7.18 (s, 1H, H1'), 7.40-7.52 (m, 6H), 7.63-7.74 (m, 5H, including NH₂), 7.91-8.04 (m, 6H), 8.96 (s, 1H, H8),


Mass spectrum: *m/z* (FAB>0) 1269 (2M+H)⁺, 635 (M+H)⁺, (FAB<0) 633 (M-H)⁻

4-amino-7-azido-(2-C-methyl- β -D-ribofuranosyl)imidazo[4,5-*d*]pyridazine or 7-amino-4-azido-1-(2-C-methyl- β -D-ribofuranosyl)imidazo[4,5-*d*]pyridazine:

¹H NMR (DMSO-*d*₆) δ ppm: 0.75 (s, 3H, CH₃), 3.65-4.15 (m, 4H), 5.30-5.55 (br, 3H, 3xOH), 6.27 (s, 1H, H1'), 7.63 (br, 2H, NH₂), 9.03 (s, 1H, H8),

Mass spectrum: *m/z* (FAB>0) 323 (M+H)⁺, (FAB<0) 321 (M-H)

Example 11: Synthesis of 4-amino-6-substituted-imidazo[4,5-*d*]- ν -triazine nucleosides



Step A : 4-amino-6-bromo-7-(β -D-ribofuranosyl)imidazo[4,5-*d*]- ν -triazine:

The 2-azaadenosine [for preparation see Patent WO 01/16149, 2001] (70 mg, 0.26 mmol) was added to a solution of sodium acetate 0.5M (1.4 mL). The solution was heated until the 2-azaadenosine was solubilized. A solution of bromine (100 μ L of Br₂ in 10 mL of water) (6.3 mL, 1.22 mmol) was added and the mixture was stirred at 20°C for 3 days. A second portion of the bromine's solution (6.3 mL, 1.22 mmol) was added and the mixture was stirred at 20°C for 3 hours. The reaction mixture was evaporated to

dryness. The crude product was purified on silica gel reverse-phase (C18) using water/acetonitrile (9/1) as eluant to give the title compound as a yellow powder.

^1H NMR (DMSO- d_6) δ ppm: 3.55 (m, 1H, H_5'), 3.71 (m, 1H, H_5'), 4.01 (m, 1H, H_4'), 4.31 (m, 1H, H_3'), 5.17 (m, 1H, H_2'), 5.19 (m, 1H, OH), 5.36 (m, 1H, OH), 5.58 (m, 1H, OH), 5.93 (d, 1H, $J=6.47$ Hz, H_1'), 8.08 (br, 2H, NH_2).

Mass spectrum: m/z (FAB >0) 349 ($\text{M}+2\text{H}$) $^+$, m/z (FAB <0) 345 ($\text{M}-2\text{H}$) $^-$.


Step B : 4-amino-6-methyl-7-(β -D-ribofuranosyl)imidazo[4,5- d]- ν -triazine:

The compound from Step A (112 mg, 0.3 mmol) was heated at reflux in hexamethyldisilazane (15 mL) for 16 hours. The mixture was evaporated to dryness to give a syrup which was dissolved in dry THF (12 mL). PPh_3 (10 mg ; 0.04 mmol), PdCl_2 (3.5 mg ; 0.02 mmol) and AlMe_3 (100 μL ; 0.94 mmol) were added. The mixture was reflux for 5 hours. The mixture was evaporated to dryness. The crude product was dissolved in methanol (30 mL) in the presence of ammonium chloride. The mixture was evaporated to dryness and the residue was purified on silica gel reverse-phase (C18) using water/acetonitrile (from 9/1 to 6/4) as eluant to give the title compound (35 mg) as a yellow powder.

^1H NMR (DMSO- d_6) δ ppm: 2.67 (s, 3H, CH_3), 3.60 (m, 1H, H_5'), 3.72 (m, 1H, H_5'), 4.03 (m, 1H, H_4'), 4.24 (m, 1H, H_3'), 4.93 (m, 1H, H_2'), 5.48 (m, 3H, OH), 5.92 (d, 1H, $J=6.82$ Hz, H_1'), 7.78 (br, 2H, NH_2).

Mass spectrum: m/z (FAB >0) (FAB >0) 283 ($\text{M}+\text{H}$) $^+$, m/z (FAB <0) 281 ($\text{M}-\text{H}$) $^-$.

Example 12: Synthesis of imidazo[4,5- d]- ν -triazin-4-one nucleosides



Step A : 7-(β -D-Ribofuranosyl)imidazo[4,5-*d*]- ν -triazin-4-one:

5 The AICAR [for preparation see *Synthesis*, 2003, No 17, 2639] (1 g, 3.87 mmol) was added to a solution of chlorhydrique acid 6N (25 mL) at -30°C. A solution of sodium nitrite 3M (4 ml, 11.62 mmol) was added and the mixture was stirred at -30° for 2 hours. A pre-cooled (-30°C) solution of ethanol (25 mL) was added. A solution of ammonia (28%) was added at -20°C to pH=7. The reaction mixture was evaporated to dryness. The crude product was purified on silica gel reverse-phase (C18) using water as eluant to give the title compound (0.81 gr) as a white powder.

10 ¹H NMR (DMSO-*d*₆) δ ppm: 3.58 (d, 1H, J=11.85Hz, H_{5'}), 3.70 (d, 1H, J=11.85Hz, H_{5'}), 4.00 (dd, 1H, J=3.92Hz, 4.02Hz, H_{4'}), 4.18 (dd, 1H, J=4.27Hz, 4.78Hz, H_{3'}), 4.54 (dd, 1H, J=4.86Hz, 5.19Hz, H_{2'}), 5.18 (br, 1H, OH), 5.35 (br, 1H, OH), 5.73 (br, 1H, OH), 6.08 (d, 1H, J=5.11Hz, H_{1'}), 8.65 (s, 1H, H₈).

Step B : 7-(2,3,5-Tri-*O*-acétyl- β -D-ribofuranosyl)imidazo[4,5-*d*]- ν -triazin-4-one:

15 The compound from Step A (1.68 gr, 6.24 mmol) was stirred in pyridine (20 mL). The anhydride acetic (2.3 ml, 25 mmol) was added and the mixture was stirred at 20°C for 16 hours. The mixture was evaporated to dryness to give a syrup which was dissolved in water. The aqueous layer was extracted by acetyl acetate. The organic layer was evaporated to dryness to give the title compound (1.5 gr) as a brown foam.


20 ¹H NMR (DMSO-*d*₆) δ ppm: 2.04 (s, 3H, COCH₃), 2.09 (s, 3H, COCH₃), 2.10 (s, 3H, COCH₃), 4.39 (m, 3H, 2xH_{5'} et H_{4'}), 5.53 (dd, 1H, J=4.39Hz, 5.4Hz, H_{3'}), 5.80 (t, 1H, J=5.4Hz, H_{2'}), 6.26 (d, 1H, J=5.4Hz, H_{1'}), 8.15 (s, 1H, H₈).

Example 13: Alternative Methods for Ribofuranosyl-Purine Analogues Synthesis

25 I. *Preparation of 4-methylamino-7-(β -D-ribofuranosyl)imidazo[4,5-*d*]- ν -triazine:*

The 4-methylamino-7-(β -D-ribofuranosyl)imidazo[4,5-*d*]- ν -triazine Va may be prepared according the following synthesis, where the starting material used is the AICAR I. The AICAR may be prepared according to the published synthesis of Y.Yamamoto and N.Kohyama, *Synthesis*, 2003, 17:2639-2646.

The other synthesis of 4-methylamino-7-(β -D-ribofuranosyl)imidazo[4,5-d]-*v*-triazine **Va** was described from 2-azainosine **II** according to the published synthesis of L.Townsend and Co, *Nucleosides, Nucleotides & Nucleic Acids*, 2000, 19(1&2):39-68.




5

II. Preparation of 4-substituted-7-(2,3-dideoxy- β -D-glycero-pentofuranosyl)-imidazo[4,5-d]-*v*-triazine derivative compounds:


The 4-substituted-7-(2,3-dideoxy- β -D-glycero-pentofuranosyl)imidazo[4,5-d]-*v*-triazine compounds **IXa**, **IXb** and **IXc** may be prepared according the following synthesis according to the published synthesis of R.Panzica and Co, *Bioorganic & Medicinal Chemistry*, 1999, 7:2373-2379.

10



III. Preparation of 4-substituted-7-(2,3-dideoxy-β-D-glycero-pent-2-ene-furanosyl)-imidazo-[4,5-d]-v-triazine derivative compounds:

5 The 4-substituted-7-(2,3-dideoxy-β-D-glycero-pent-2-ene-furanosyl)imidazo[4,5-d]-v-triazine derivative compounds **XIa**, **XIb** and **XIc** may be prepared according the following synthesis:



Example 14: Phosphorylation Assay of Nucleoside to Active Triphosphate

To determine the cellular metabolism of the compounds, HepG2 cells are obtained from the American Type Culture Collection (Rockville, MD), and are grown in 225 cm² tissue culture flasks in minimal essential medium supplemented with non-essential amino acids, 1% penicillin-streptomycin. The medium is renewed every three days, and the cells are subcultured once a week. After detachment of the adherent monolayer with a 10 minute exposure to 30 mL of trypsin-EDTA and three consecutive washes with medium, confluent HepG2 cells are seeded at a density of 2.5×10^6 cells per well in a 6-well plate and exposed to 10 μ M of [³H] labeled active compound (500 dpm/pmol) for the specified time periods. The cells are maintained at 37°C under a 5% CO₂ atmosphere. At the selected time points, the cells are washed three times with ice-cold phosphate-buffered saline (PS). Intracellular active compound and its respective metabolites are extracted by incubating the cell pellet overnight at -20°C with 60% methanol followed by extraction with an additional 20 μ L of cold methanol for one hour in an ice bath. The extracts are then combined, dried under gentle filtered air flow and stored at -20°C until HPLC analysis.

Example 15: Bioavailability Assay in Cynomolgus Monkeys

Within 1 week prior to the study initiation, the cynomolgus monkey is surgically implanted with a chronic venous catheter and subcutaneous venous access port (VAP) to facilitate blood collection and underwent a physical examination including hematology and serum chemistry evaluations and the body weight was recorded. Each monkey (six total) receives approximately 250 μ Ci of ³H activity with each dose of active compound at a dose level of 10 mg/kg at a dose concentration of 5 mg/mL, either via an intravenous route (3 monkeys, IV), or via oral gavage (3 monkeys, PO). Each dosing syringe is weighed before dosing to gravimetrically determine the quantity of formulation administered. Urine samples are collected via pan catch at the designated intervals (approximately 18-0 hours pre-dose, 0-4, 4-8 and 8-12 hours post-dosage) and processed. blood samples are collected as well (pre-dose, 0.25, 0.5, 1, 2, 3, 6, 8, 12 and 24 hours post-dosage) via the chronic venous catheter and VAP or from a peripheral vessel if the chronic venous catheter procedure should not be possible. The blood and urine samples

are analyzed for the maximum concentration (C_{max}), time when the maximum concentration is achieved (T_{max}), area under the curve (AUC), half life of the dosage concentration ($T_{1/2}$), clearance (CL), steady state volume and distribution (V_{ss}) and bioavailability (F).

5

Example 16: Bone Marrow Toxicity Assay

Human one marrow cells are collected from normal healthy volunteers and the mononuclear population are separated by Ficoll-Hypaque gradient centrifugation as described previously by Sommadossi J-P, Carlisle R. "Toxicity of 3'-azido-3'-deoxythymidine and 9-(1,3-dihydroxy-2-propoxymethyl)guanine for normal human hematopoietic progenitor cells *in vitro*" *Antimicrobial Agents and Chemotherapy* 1987; 31:452-454; and Sommadossi J-P, Schinazi RF, Chu CK, Xie M-Y. "Comparison of cytotoxicity of the (-)- and (+)-enantiomer of 2',3'-dideoxy-3'-thiacytidine in normal human one marrow progenitor cells" *Biochemical Pharmacology* 1992; 44:1921-1925.

10

15

The culture assays for CFU-GM and FU-E are performed using a bilayer soft agar or methylcellulose method. Drugs are diluted in tissue culture medium and filtered. After 14 to 18 days at 37°C in a humidified atmosphere of 5% CO₂ in air, colonies of greater than 50 cells are counted using an inverted microscope. The results are presented as the percent inhibition of colony formation in the presence of drug compared to solvent control

20

cultures.

Example 17: Mitochondria Toxicity Assay

HepG2 cells are cultured in 12-well plates as described above and exposed to various concentrations of drugs as taught by Pan-Zhou X-R, Cui L, Zhou X-J, Sommadossi J-P, Darley-Usmer VM. "Differential effects of antiretroviral nucleoside analogs on mitochondrial function in HepG2 cells" *Antimicro Agents Chemother* 2000; 44:496-503. Lactic acid levels in the culture medium after 4 day drug exposure are measured using a Boehringer lactic acid assay kit. Lactic acid levels are normalized by cell number as measured by hemocytometer count.

25

30

Example 18: Cytotoxicity Assay

Cells are seeded at a rate of between 5×10^3 and 5×10^4 /well into 96-well plates in growth medium overnight at 37°C in a humidified CO₂ (5%) atmosphere. New growth medium containing serial dilutions of the drugs is then added. After incubation for 4 days, cultures are fixed in 50% TCA and stained with sulforhodamine. The optical density was read at 550 nm. The cytotoxic concentration was expressed as the concentration required to reduce the cell number by 50% (CC₅₀). The preliminary results are tabulated in the Table 3 below.


Table 3: MDK versus Human Hepatoma

Compound	CC ₅₀ , μ M
	MDK
β -D-4'-CH ₃ -riboG	>250
β -D-4'-CH ₃ -ribo-4-thioU	>250
β -D-4'-CH ₃ -riboC	>250
β -D-4'-CH ₃ -ribo-5-fluoroU	>167
β -D-4'-CH ₃ -riboT	>250
β -D-4'-CH ₃ -riboA	>250

This invention has been described with reference to its preferred embodiments. Variations and modifications of the invention will be obvious to those skilled in the art from the foregoing detailed description of the invention. It is intended that all of these variations and modifications be included within the scope of this invention.

WE CLAIM:

1. A method of treating a host infected with a flavivirus or pestivirus, comprising administering an effective amount of an anti-pestivirus or anti-flavivirus biologically active ribofuranonucleoside of Formula (I):



(I)

or a pharmacologically acceptable salt or prodrug thereof, wherein:

R is H, mono-, di-, or triphosphate, a stabilized phosphate, or phosphonate;

X is O, S[O]_n, CH₂, CHOH, CH-alkyl, CH-alkenyl, CH-alkynyl, C-dialkyl, CH-O-alkyl, CH-O-alkenyl, CH-O-alkynyl, CH-S-alkyl, CH-S-alkenyl, CH-S-alkynyl, NH, N-alkyl, N-alkenyl, N-alkynyl, S(O)N-alkyl, S(O)N-alkenyl, S(O)N-alkynyl, SCH-halogen, or C-(halogen)₂, wherein alkyl, alkenyl or alkynyl may optionally be substituted;

n is 0-2;

such than when X is CH₂, CHOH, CH-alkyl, CH-alkenyl, CH-alkynyl, C-dialkyl, CH-O-alkyl, CH-O-alkenyl, CH-O-alkynyl, CH-S-alkyl, CH-S-alkenyl, CH-S-alkynyl, CH-halogen, or C-(halogen)₂,

then each R¹ and R^{1'} is independently H, OH, optionally substituted alkyl, lower alkyl, azido, cyano, optionally substituted alkenyl or alkynyl, -C(O)O-(alkyl), -C(O)O(lower alkyl), -C(O)O-(alkenyl), -C(O)O-(alkynyl), -O(acyl), -O(lower acyl), -O(alkyl), -O(lower alkyl), -O(alkenyl), -O(alkynyl), halogen, halogenated alkyl, -NO₂, -NH₂, -NH(lower alkyl), -N(lower alkyl)₂, -NH(acyl), -N(acyl)₂, -C(O)NH₂, -C(O)NH(alkyl), -C(O)N(alkyl)₂, S(O)N-

alkyl, S(O)N-alkenyl, S(O)N-alkynyl, SCH-halogen, wherein alkyl, alkenyl, and/or alkynyl may optionally be substituted; and


such that when X is O, S[O]_n, NH, N-alkyl, N-alkenyl, N-alkynyl, S(O)N-alkyl, S(O)N-alkenyl, S(O)N-alkynyl, or SCH-halogen,

5 then each R¹ and R^{1'} is independently H, optionally substituted alkyl, lower alkyl, azido, cyano, optionally substituted alkenyl or alkynyl, -C(O)O-(alkyl), -C(O)O(lower alkyl), -C(O)O-(alkenyl), -C(O)O-(alkynyl), halogenated alkyl, -C(O)NH₂, -C(O)NH(alkyl), -C(O)N(alkyl)₂, -C(H)=N-NH₂, C(S)NH₂, C(S)NH(alkyl), or C(S)N(alkyl)₂, wherein alkyl, alkenyl and/or alkynyl may
10 optionally be substituted;

each R² and R³ independently is OH, NH₂, SH, F, Cl, Br, I, CN, NO₂, -C(O)NH₂, -C(O)NH(alkyl), and -C(O)N(alkyl)₂, N₃, optionally substituted alkyl, lower
15 alkyl, optionally substituted alkenyl or alkynyl, halogenated alkyl, -C(O)O-(alkyl), -C(O)O(lower alkyl), -C(O)O-(alkenyl), -C(O)O-(alkynyl), -O(acyl), -O(alkyl), -O(alkenyl), -O(alkynyl), -OC(O)NH₂, NC, C(O)OH, SCN, OCN, -S(alkyl), -S(alkenyl), -S(alkynyl), -NH(alkyl), -N(alkyl)₂, -NH(alkenyl), -NH(alkynyl), an amino acid residue or derivative, a prodrug or leaving group that provides OH *in vivo*, or an optionally substituted 3-7 membered heterocyclic ring having O, S and/or N independently as a heteroatom taken
20 alone or in combination;

each R^{2'} and R^{3'} independently is H; optionally substituted alkyl, alkenyl, or
alkynyl; -C(O)O(alkyl), -C(O)O(lower alkyl), -C(O)O(alkenyl), -C(O)O(alkynyl), -C(O)NH₂, -C(O)NH(alkyl), -C(O)N(alkyl)₂, -O(acyl), -O(lower acyl), -O(alkyl), -O(lower alkyl), -O(alkenyl), halogen, halogenated
25 alkyl and particularly CF₃, azido, cyano, NO₂, -S(alkyl), -S(alkenyl), -S(alkynyl), NH₂, -NH(alkyl), -N(alkyl)₂, -NH(alkenyl), -NH(alkynyl), -NH(acyl), or -N(acyl)₂, and R₃ at 3'-C may also be OH; and

Base is selected from the group consisting of:



5 wherein

each R' and R'' independently is H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, halogen, halogenated alkyl, OH, CN, N₃, carboxy, C₁₋₄alkoxycarbonyl, NH₂, C₁₋₄ alkylamino, di(C₁₋₄ alkyl)amino, C₁₋₆ alkoxy, C₁₋₆ alkylsulfonyl, (C₁₋₄ alkyl)₀₋₂ aminomethyl;


10 each W is Cl, Br, I, F, halogenated alkyl, alkoxy, OH, SH, O-alkyl, S-alkyl, O-alkenyl, O-alkynyl, S-alkenyl, S-alkynyl, -OC(O)NR⁴R⁴, O-acyl, S-acyl, CN, SCN, OCN, NO₂, N₃, NH₂, NH(alkyl), N(alkyl)₂, NH-cycloalkyl, NH-acyl, NH=NH, CONH₂, CONH(alkyl), or CON(alkyl)₂; and

each R⁴ is independently H, acyl, or C₁₋₆ alkyl;

15 each Z is O, S, NH, N-OH, N-NH₂, NH(alkyl), N(alkyl)₂, N-cycloalkyl, alkoxy, CN, SCN, OCN, SH, NO₂, NH₂, N₃, NH=NH, NH(alkyl), N(alkyl)₂, CONH₂, CONH(alkyl), or CON(alkyl)₂;

with the caveat that when X is S, then the compound is not 5-(4-amino-imidazo[4,5-d][1,2,3]triazin-7-yl)-2-hydroxymethyl-tetrahydro-thiophen-3-ol or 7-(4-hydroxy-5-hydroxy-methyl-tetrahydro-thiophen-2-yl)-3,7-dihydro-imidazo[4,5-d][1,2,3]triazin-4-one.

- 5 2. A method of treating a host infected with a flavivirus or pestivirus, comprising administering an effective amount of an anti-pestivirus or anti-flavivirus biologically active ribofuranonucleoside of Formula (II):



(II)

10 or a pharmacologically acceptable salt or prodrug thereof, wherein:

X^* is CY^3 ;

Y^3 is hydrogen, alkyl, bromo, chloro, fluoro, iodo, azido, cyano, alkenyl, alkynyl, $-C(O)O(alkyl)$, $-C(O)O(lower\ alkyl)$, CF_3 , $-CONH_2$, $-CONH(alkyl)$, or $-CON(alkyl)_2$;

15 R is H, mono-, di-, or triphosphate, a stabilized phosphate, or phosphonate;

R^1 is H, OH, optionally substituted alkyl, lower alkyl, azido, cyano, optionally substituted alkenyl or alkynyl, $-C(O)O-(alkyl)$, $-C(O)O(lower\ alkyl)$, $-C(O)O-(alkenyl)$, $-C(O)O-(alkynyl)$, $-O(acyl)$, $-O(lower\ acyl)$, $-O(alkyl)$, $-O(lower\ alkyl)$, $-O(alkenyl)$, $-O(alkynyl)$, halogen, halogenated alkyl, $-NO_2$, $-NH_2$, $-NH(lower\ alkyl)$, $-N(lower\ alkyl)_2$, $-NH(acyl)$, $-N(acyl)_2$, $-C(O)NH_2$, $-C(O)NH(alkyl)$, or $-C(O)N(alkyl)_2$, wherein an optional substitution on alkyl, alkenyl, and/or alkynyl may be one or more halogen, hydroxy, alkoxy or alkylthio groups taken in any combination;

20


each R^2 and R^3 independently is OH, NH_2 , F, Cl, Br, I, CN, NO_2 , $-C(O)NH_2$, $-C(O)NH(alkyl)$, $-C(O)N(alkyl)_2$, N_3 , optionally substituted alkyl, lower

25

alkyl, optionally substituted alkenyl or alkynyl, halogenated alkyl, -C(O)O-
 (alkyl), -C(O)O(lower alkyl), -C(O)O-(alkenyl), -C(O)O-(alkynyl), an amino
 acid residue or derivative, a prodrug or leaving group that provides OH *in*
in vivo, or an optionally substituted 3-7 membered heterocyclic ring having O, S
 and/or N independently as a heteroatom taken alone or in combination;

each R^{2'} and R^{3'} independently is H; optionally substituted alkyl, alkenyl, or
 alkynyl; -C(O)O(alkyl), -C(O)O(lower alkyl), -C(O)O(alkenyl),
 -C(O)O(alkynyl), -C(O)NH₂, -C(O)NH(alkyl), and -C(O)N(alkyl)₂, -O(acyl),
 -O(lower acyl), -O(alkyl), -O(lower alkyl), -O(alkenyl), halogen, halogenated
 alkyl and particularly CF₃, azido, cyano, NO₂, -S(alkyl), -S(alkenyl),
 -S(alkynyl), NH₂, -NH(alkyl), -N(alkyl)₂, -NH(alkenyl), -NH(alkynyl),
 -NH(acyl), or -N(acyl)₂, and R₃ at 3'-C may also be OH; and

Base is selected from the group consisting of:



wherein


each R' and R'' independently is H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, halogen, halogenated alkyl, OH, CN, N₃, carboxy, C₁₋₄alkoxycarbonyl, NH₂, C₁₋₄ alkylamino, di(C₁₋₄ alkyl)amino, C₁₋₆ alkoxy, C₁₋₆ alkylsulfonyl, (C₁₋₄ alkyl)_{0.2} aminomethyl;

5 each W is Cl, Br, I, F, halogenated alkyl, alkoxy, OH, SH, O-alkyl, S-alkyl, O-alkenyl, O-alkynyl, S-alkenyl, S-alkynyl, -OC(O)NR⁴R⁴, O-acyl, S-acyl, CN, SCN, OCN, NO₂, N₃, NH₂, NH(alkyl), N(alkyl)₂, NH-cycloalkyl, NH-acyl, NH=NH, CONH₂, CONH(alkyl), or CON(alkyl)₂; and

each R⁴ is independently H, acyl, or C₁₋₆ alkyl;

10 each Z is O, S, NH, N-OH, N-NH₂, NH(alkyl), N(alkyl)₂, N-cycloalkyl, alkoxy, CN, SCN, OCN, SH, NO₂, NH₂, N₃, NH=NH, NH(alkyl), N(alkyl)₂, CONH₂, CONH(alkyl), or CON(alkyl)₂.

3. A method of treating a host infected with a flavivirus or pestivirus, comprising administering an effective amount of an anti-pestivirus or anti-flavivirus
15 biologically active ribofuranonucleoside of Formula (III):



(III)

or a pharmacologically acceptable salt or prodrug thereof, wherein:


20 each R, R^{2*}, and R^{3*} independently is H, mono-, di-, or triphosphate, a stabilized phosphate, or phosphonate; optionally substituted alkyl, lower alkyl, optionally substituted alkenyl or alkynyl, acyl, -C(O)-(alkyl), -C(O)(lower alkyl), -C(O)-(alkenyl), -C(O)-(alkynyl), lipid, phospholipid, carbohydrate, peptide, cholesterol, an amino acid residue or derivative, or other
25 pharmaceutically acceptable leaving group that is capable of providing H or phosphate when administered *in vivo*;

X is O, S[O]_n, CH₂, CHOH, CH-alkyl, CH-alkenyl, CH-alkynyl, C-dialkyl, CH-O-alkyl, CH-O-alkenyl, CH-O-alkynyl, CH-S-alkyl, CH-S-alkenyl, CH-S-alkynyl, NH, N-alkyl, N-alkenyl, N-alkynyl, S(O)N-alkyl, S(O)N-alkenyl, S(O)N-alkynyl, SCH-halogen, or C-(halogen)₂, wherein alkyl, alkenyl or alkynyl optionally may be substituted;

n is 0-2;

each R^{2'} independently is H; optionally substituted alkyl, alkenyl, or alkynyl; -C(O)O(alkyl), -C(O)O(lower alkyl), -C(O)O(alkenyl), -C(O)O(alkynyl), -C(O)NH₂, -C(O)NH(alkyl), -C(O)N(alkyl)₂, -OH, -O(acyl), -O(lower acyl), -O(alkyl), -O(lower alkyl), -O(alkenyl), halogen, halogenated alkyl and particularly CF₃, azido, cyano, NO₂, -S(alkyl), -S(alkenyl), -S(alkynyl), NH₂, -NH(alkyl), -N(alkyl)₂, -NH(alkenyl), -NH(alkynyl), -NH(acyl), or -N(acyl)₂; and


Base is selected from the group consisting of:



wherein

- each R' and R'' independently is H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, halogen, halogenated alkyl, OH, CN, N₃, carboxy, C₁₋₄alkoxycarbonyl, NH₂, C₁₋₄ alkylamino, di(C₁₋₄ alkyl)amino, C₁₋₆ alkoxy, C₁₋₆ alkylsulfonyl, (C₁₋₄ alkyl)₀₋₂ aminomethyl;
- 5 each W is Cl, Br, I, F, halogenated alkyl, alkoxy, OH, SH, O-alkyl, S-alkyl, O-alkenyl, O-alkynyl, S-alkenyl, S-alkynyl, -OC(O)NR⁴R⁴, O-acyl, S-acyl, CN, SCN, OCN, NO₂, N₃, NH₂, NH(alkyl), N(alkyl)₂, NH-cycloalkyl, NH-acyl, NH=NH, CONH₂, CONH(alkyl), or CON(alkyl)₂; and
- each R⁴ is independently H, acyl, or C₁₋₆ alkyl;
- 10 each Z is O, S, NH, N-OH, N-NH₂, NH(alkyl), N(alkyl)₂, N-cycloalkyl, alkoxy, CN, SCN, OCN, SH, NO₂, NH₂, N₃, NH=NH, NH(alkyl), N(alkyl)₂, CONH₂, CONH(alkyl), or CON(alkyl)₂.
4. The method of claim 3, wherein R^{2'} is an optionally substituted alkyl, alkenyl, or alkynyl; halogen, halogenated alkyl, CH₃, CF₃, azido, or cyano.
- 15 5. The method of claim 3, wherein R^{2'} is an optionally substituted alkyl, alkenyl, or alkynyl; halogen, halogenated alkyl, CH₃, or CF₃.
6. The method of claim 3, wherein R^{2'} is CH₃ or CF₃.
7. The method of claim 3, wherein each R, R^{2*}, and R^{3*} is independently H, mono-, di-, or triphosphate, a stabilized phosphate, or phosphonate.
- 20 8. The method of claim 3, wherein each R, R^{2*}, and R^{3*} is independently H.
9. The method of claim 3, wherein each R, R^{2*}, and R^{3*} is independently H, acyl, or an amino acid acyl residue.
10. The method of claim 3, wherein X is O or S.
11. The method of claim 3, wherein X is O.

12. A method of treating a host infected with a flavivirus or pestivirus, comprising administering an effective amount of an anti-pestivirus or anti-flavivirus biologically active ribofuranonucleoside of Formula (IV):



(IV)

or a pharmacologically acceptable salt or prodrug thereof, wherein:


each R, R^{2*}, and R^{3*} independently is H, mono-, di-, or triphosphate, a stabilized phosphate, or phosphonate; optionally substituted alkyl, lower alkyl, optionally substituted alkenyl or alkynyl, acyl, -C(O)-(alkyl), -C(O)(lower alkyl), -C(O)-(alkenyl), -C(O)-(alkynyl), lipid, phospholipid, carbohydrate, peptide, cholesterol, an amino acid residue or derivative, or other pharmaceutically acceptable leaving group that is capable of providing H or phosphate when administered *in vivo*;

X is O, S[O]_n, CH₂, CHOH, CH-alkyl, CH-alkenyl, CH-alkynyl, C-dialkyl, CH-O-alkyl, CH-O-alkenyl, CH-O-alkynyl, CH-S-alkyl, CH-S-alkenyl, CH-S-alkynyl, NH, N-alkyl, N-alkenyl, N-alkynyl, S(O)N-alkyl, S(O)N-alkenyl, S(O)N-alkynyl, SCH-halogen, or C-(halogen)₂, wherein alkyl, alkenyl or alkynyl optionally may be substituted;

n is 0-2;

each R^{3'} independently is H; optionally substituted alkyl, alkenyl, or alkynyl; -C(O)O(alkyl), -C(O)O(lower alkyl), -C(O)O(alkenyl), -C(O)O(alkynyl), -C(O)NH₂, -C(O)NH(alkyl), -C(O)N(alkyl)₂, -OH, -O(acyl), -O(lower acyl), -O(alkyl), -O(lower alkyl), -O(alkenyl), halogen, halogenated alkyl and particularly CF₃, azido, cyano, NO₂, -S(alkyl), -S(alkenyl), -S(alkynyl), NH₂, -NH(alkyl), -N(alkyl)₂, -NH(alkenyl), -NH(alkynyl), -NH(acyl), or -N(acyl)₂; and

Base is selected from the group consisting of:



5

wherein

each R' and R'' independently is H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, halogen, halogenated alkyl, OH, CN, N₃, carboxy, C₁₋₄alkoxycarbonyl, NH₂, C₁₋₄ alkylamino, di(C₁₋₄ alkyl)amino, C₁₋₆ alkoxy, C₁₋₆ alkylsulfonyl, (C₁₋₄ alkyl)₀₋₂ aminomethyl;

10

each W is Cl, Br, I, F, halogenated alkyl, alkoxy, OH, SH, O-alkyl, S-alkyl, O-alkenyl, O-alkynyl, S-alkenyl, S-alkynyl, -OC(O)NR⁴R⁴, O-acyl, S-acyl, CN, SCN, OCN, NO₂, N₃, NH₂, NH(alkyl), N(alkyl)₂, NH-cycloalkyl, NH-acyl, NH=NH, CONH₂, CONH(alkyl), or CON(alkyl)₂; and


each R⁴ is independently H, acyl, or C₁₋₆ alkyl;

15

each Z is O, S, NH, N-OH, N-NH₂, NH(alkyl), N(alkyl)₂, N-cycloalkyl, alkoxy, CN, SCN, OCN, SH, NO₂, NH₂, N₃, NH=NH, NH(alkyl), N(alkyl)₂, CONH₂, CONH(alkyl), or CON(alkyl)₂.

13. The method of claim 12, wherein R^{3'} is an optionally substituted alkyl, alkenyl, or alkynyl; halogen, halogenated alkyl, CH₃, CF₃, azido, or cyano.
14. The method of claim 12, wherein R^{3'} is an optionally substituted alkyl, alkenyl, or alkynyl; halogen, halogenated alkyl, CH₃, or CF₃.
- 5 15. The method of claim 12, wherein R^{3'} is CH₃ or CF₃.
16. The method of claim 12, wherein each R, R^{2*}, and R^{3*} is independently H, mono-, di-, or triphosphate, a stabilized phosphate, or phosphonate.
17. The method of claim 12, wherein each R, R^{2*}, and R^{3*} is independently H.
18. The method of claim 12, wherein each R, R^{2*}, and R^{3*} is independently H, acyl,
10 or an amino acid acyl residue.
19. The method of claim 12, wherein X is O or S.
20. The method of claim 12, wherein X is O.
21. The method of any one of claims 1-3 and 12 wherein the host is a mammal.
22. The method of claim 21, wherein the mammal is a human.
- 15 23. The method of any one of claims 1-3 and 12, further comprising administering an antivirally effective amount of the compound, or a pharmaceutically acceptable salt or prodrug thereof, in combination or alternation with one or more additional antivirally effective agents.
- 20 24. The method of claim 23 wherein the additional antivirally effective agent is selected from the group consisting of an interferon, ribavirin, an interleukin, an NS3 protease inhibitor, a cysteine protease inhibitor, phenanthrenequinone, a thiazolidine derivative, a thiazolidine and a benzanilide, a helicase inhibitor, a polymerase inhibitor, a nucleotide analogue, gliotoxin, cerulenin, an antisense phosphorothioate oligodeoxynucleotide, an inhibitor of IRES-dependent
25 translation, and a ribozyme.
25. The method of claim 24, wherein the additional antivirally effective agent is an interferon.

26. The method of claim 25 wherein the additional antivirally effective agent is selected from the group consisting of pegylated interferon alpha 2a, interferon alphacon-1, natural interferon, albuferon, interferon beta-1a, omega interferon, interferon alpha, interferon gamma, interferon tau, interferon delta and interferon gamma-1b.
27. The method of any one of claims 1-3 and 12, wherein the compound is in the form of a dosage unit.
28. The method of claim 27 wherein the dosage unit contains 50 to 1000 mg of the compound.
29. The method of claim 28, wherein the said dosage unit is a tablet or capsule.
30. The method of any one of claims 1-3 and 12, wherein the compound is in substantially pure form.
31. The method of claim 30 wherein the compound is at least 90% by weight of the β -D-isomer.
32. The method of claim 30 wherein the compound is at least 95% by weight of the β -D-isomer.
33. The method of claim 30 wherein the compound is at least 90% by weight of the β -L-isomer.
34. The method of claim 30 wherein the compound is at least 95% by weight of the β -L-isomer.
35. A compound of the general structure of Formula (I):



(I)

or a pharmacologically acceptable salt or prodrug thereof, wherein:

R is H, mono-, di-, or triphosphate, a stabilized phosphate, or phosphonate;

X is O, S[O]_n, CH₂, CHOH, CH-alkyl, CH-alkenyl, CH-alkynyl, C-dialkyl, CH-O-alkyl, CH-O-alkenyl, CH-O-alkynyl, CH-S-alkyl, CH-S-alkenyl, CH-S-alkynyl, NH, N-alkyl, N-alkenyl, N-alkynyl, S(O)N-alkyl, S(O)N-alkenyl, S(O)N-alkynyl, SCH-halogen, or C-(halogen)₂, wherein alkyl, alkenyl or alkynyl may optionally be substituted;

n is 0-2;

such that when X is CH₂, CHOH, CH-alkyl, CH-alkenyl, CH-alkynyl, C-dialkyl, CH-O-alkyl, CH-O-alkenyl, CH-O-alkynyl, CH-S-alkyl, CH-S-alkenyl, CH-S-alkynyl, CH-halogen, or C-(halogen)₂,

then each R¹ and R^{1'} is independently H, OH, optionally substituted alkyl, lower alkyl, azido, cyano, optionally substituted alkenyl or alkynyl, -C(O)O-(alkyl), -C(O)O(lower alkyl), -C(O)O-(alkenyl), -C(O)O-(alkynyl), -O(acyl), -O(lower acyl), -O(alkyl), -O(lower alkyl), -O(alkenyl), -O(alkynyl), halogen, halogenated alkyl, -NO₂, -NH₂, -NH(lower alkyl), -N(lower alkyl)₂, -NH(acyl), -N(acyl)₂, -C(O)NH₂, -C(O)NH(alkyl), -C(O)N(alkyl)₂, S(O)N-alkyl, S(O)N-alkenyl, S(O)N-alkynyl, SCH-halogen, wherein alkyl, alkenyl, and/or alkynyl may optionally be substituted; and

such that when X is O, S[O]_n, NH, N-alkyl, N-alkenyl, N-alkynyl, S(O)N-alkyl, S(O)N-alkenyl, S(O)N-alkynyl, or SCH-halogen,

then each R¹ and R^{1'} is independently H, optionally substituted alkyl, lower alkyl, azido, cyano, optionally substituted alkenyl or alkynyl, -C(O)O-(alkyl), -C(O)O(lower alkyl), -C(O)O-(alkenyl), -C(O)O-(alkynyl), halogenated alkyl, -C(O)NH₂, -C(O)NH(alkyl), -C(O)N(alkyl)₂, -C(H)=N-NH₂, C(S)NH₂, C(S)NH(alkyl), or C(S)N(alkyl)₂, wherein alkyl, alkenyl and/or alkynyl may optionally be substituted;

each R² and R³ independently is OH, NH₂, SH, F, Cl, Br, I, CN, NO₂, -C(O)NH₂, -C(O)NH(alkyl), and -C(O)N(alkyl)₂, N₃, optionally substituted alkyl, lower alkyl, optionally substituted alkenyl or alkynyl, halogenated alkyl, -C(O)O-(alkyl), -C(O)O(lower alkyl), -C(O)O-(alkenyl), -C(O)O-(alkynyl), -O(acyl),

5


-O(alkyl), -O(alkenyl), -O(alkynyl), -OC(O)NH₂, NC, C(O)OH, SCN, OCN, -S(alkyl), -S(alkenyl), -S(alkynyl), -NH(alkyl), -N(alkyl)₂, -NH(alkenyl), -NH(alkynyl), an amino acid residue or derivative, a prodrug or leaving group that provides OH *in vivo*, or an optionally substituted 3-7 membered heterocyclic ring having O, S and/or N independently as a heteroatom taken alone or in combination;

10

each R^{2'} and R^{3'} independently is H; optionally substituted alkyl, alkenyl, or alkynyl; -C(O)O(alkyl), -C(O)O(lower alkyl), -C(O)O(alkenyl), -C(O)O(alkynyl), -C(O)NH₂, -C(O)NH(alkyl), -C(O)N(alkyl)₂, -O(acyl), -O(lower acyl), -O(alkyl), -O(lower alkyl), -O(alkenyl), halogen, halogenated alkyl and particularly CF₃, azido, cyano, NO₂, -S(alkyl), -S(alkenyl), -S(alkynyl), NH₂, -NH(alkyl), -N(alkyl)₂, -NH(alkenyl), -NH(alkynyl), -NH(acyl), or -N(acyl)₂, and R₃ at 3'-C may also be OH; and

Base is selected from the group consisting of:

15



wherein

each R' and R'' independently is H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, halogen, halogenated alkyl, OH, CN, N₃, carboxy, C₁₋₄alkoxycarbonyl, NH₂, C₁₋₄ alkylamino, di(C₁₋₄ alkyl)amino, C₁₋₆ alkoxy, C₁₋₆ alkylsulfonyl, (C₁₋₄ alkyl)₀₋₂ aminomethyl;


5 each W is Cl, Br, I, F, halogenated alkyl, alkoxy, OH, SH, O-alkyl, S-alkyl, O-alkenyl, O-alkynyl, S-alkenyl, S-alkynyl, -OC(O)NR⁴R⁴, O-acyl, S-acyl, CN, SCN, OCN, NO₂, N₃, NH₂, NH(alkyl), N(alkyl)₂, NH-cycloalkyl, NH-acyl, NH=NH, CONH₂, CONH(alkyl), or CON(alkyl)₂; and

each R⁴ is independently H, acyl, or C₁₋₆ alkyl;

10 each Z is O, S, NH, N-OH, N-NH₂, NH(alkyl), N(alkyl)₂, N-cycloalkyl, alkoxy, CN, SCN, OCN, SH, NO₂, NH₂, N₃, NH=NH, NH(alkyl), N(alkyl)₂, CONH₂, CONH(alkyl), or CON(alkyl)₂;

with the caveat that when X is S, then the compound is not 5-(4-amino-imidazo[4,5-d][1,2,3]triazin-7-yl)-2-hydroxymethyl-tetrahydro-thiophen-3-ol
15 or 7-(4-hydroxy-5-hydroxy-methyl-tetrahydro-thiophen-2-yl)-3,7-dihydro-imidazo[4,5-d][1,2,3]triazin-4-one.

36. A compound of the general structure of Formula (II):



(II)

20 or a pharmacologically acceptable salt or prodrug thereof, wherein:

X* is CY³;

Y³ is hydrogen, alkyl, bromo, chloro, fluoro, iodo, azido, cyano, alkenyl, alkynyl, -C(O)O(alkyl), -C(O)O(lower alkyl), CF₃, -CONH₂, -CONH(alkyl), or -CON(alkyl)₂;


R is H, mono-, di-, or triphosphate, a stabilized phosphate, or phosphonate;

R¹ is H, OH, optionally substituted alkyl, lower alkyl, azido, cyano, optionally substituted alkenyl or alkynyl, -C(O)O-(alkyl), -C(O)O(lower alkyl), -C(O)O-(alkenyl), -C(O)O-(alkynyl), -O(acyl), -O(lower acyl), -O(alkyl), -O(lower alkyl), -O(alkenyl), -O(alkynyl), halogen, halogenated alkyl, -NO₂, -NH₂, -NH(lower alkyl), -N(lower alkyl)₂, -NH(acyl), -N(acyl)₂, -C(O)NH₂, -C(O)NH(alkyl), or -C(O)N(alkyl)₂, wherein an optional substitution on alkyl, alkenyl, and/or alkynyl may be one or more halogen, hydroxy, alkoxy or alkylthio groups taken in any combination;


each R² and R³ independently is OH, NH₂, F, Cl, Br, I, CN, NO₂, -C(O)NH₂, -C(O)NH(alkyl), -C(O)N(alkyl)₂, N₃, optionally substituted alkyl, lower alkyl, optionally substituted alkenyl or alkynyl, halogenated alkyl, -C(O)O-(alkyl), -C(O)O(lower alkyl), -C(O)O-(alkenyl), -C(O)O-(alkynyl), an amino acid residue or derivative, a prodrug or leaving group that provides OH *in vivo*, or an optionally substituted 3-7 membered heterocyclic ring having O, S and/or N independently as a heteroatom taken alone or in combination;

each R^{2'} and R^{3'} independently is H; optionally substituted alkyl, alkenyl, or alkynyl; -C(O)O(alkyl), -C(O)O(lower alkyl), -C(O)O(alkenyl), -C(O)O(alkynyl), -C(O)NH₂, -C(O)NH(alkyl), and -C(O)N(alkyl)₂, -O(acyl), -O(lower acyl), -O(alkyl), -O(lower alkyl), -O(alkenyl), halogen, halogenated alkyl and particularly CF₃, azido, cyano, NO₂, -S(alkyl), -S(alkenyl), -S(alkynyl), NH₂, -NH(alkyl), -N(alkyl)₂, -NH(alkenyl), -NH(alkynyl), -NH(acyl), or -N(acyl)₂, and R₃ at 3'-C may also be OH; and


Base is selected from the group consisting of:




(i)




(ii)



(iii)



(iv)



wherein


each R' and R'' independently is H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, halogen, halogenated alkyl, OH, CN, N₃, carboxy, C₁₋₄alkoxycarbonyl, NH₂, C₁₋₄ alkylamino, di(C₁₋₄ alkyl)amino, C₁₋₆ alkoxy, C₁₋₆ alkylsulfonyl, (C₁₋₄ alkyl)₀₋₂ aminomethyl;

each W is Cl, Br, I, F, halogenated alkyl, alkoxy, OH, SH, O-alkyl, S-alkyl, O-alkenyl, O-alkynyl, S-alkenyl, S-alkynyl, -OC(O)NR⁴R⁴, O-acyl, S-acyl, CN, SCN, OCN, NO₂, N₃, NH₂, NH(alkyl), N(alkyl)₂, NH-cycloalkyl, NH-acyl, NH=NH, CONH₂, CONH(alkyl), or CON(alkyl)₂; and

each R⁴ is independently H, acyl, or C₁₋₆ alkyl;

each Z is O, S, NH, N-OH, N-NH₂, NH(alkyl), N(alkyl)₂, N-cycloalkyl, alkoxy, CN, SCN, OCN, SH, NO₂, NH₂, N₃, NH=NH, NH(alkyl), N(alkyl)₂, CONH₂, CONH(alkyl), or CON(alkyl)₂.

37. A compound of the general structure of Formula (III):



(III)

or a pharmacologically acceptable salt or prodrug thereof, wherein:


5 each R, R^{2*}, and R^{3*} independently is H, mono-, di-, or triphosphate, a stabilized phosphate, or phosphonate; optionally substituted alkyl, lower alkyl, optionally substituted alkenyl or alkynyl, acyl, -C(O)-(alkyl), -C(O)(lower alkyl), -C(O)-(alkenyl), -C(O)-(alkynyl), lipid, phospholipid, carbohydrate, peptide, cholesterol, an amino acid residue or derivative, or other
10 pharmaceutically acceptable leaving group that is capable of providing H or phosphate when administered *in vivo*;

X is O, S[O]_n, CH₂, CHOH, CH-alkyl, CH-alkenyl, CH-alkynyl, C-dialkyl, CH-O-alkyl, CH-O-alkenyl, CH-O-alkynyl, CH-S-alkyl, CH-S-alkenyl, CH-S-alkynyl, NH, N-alkyl, N-alkenyl, N-alkynyl, S(O)N-alkyl, S(O)N-alkenyl,
15 S(O)N-alkynyl, SCH-halogen, or C-(halogen)₂, wherein alkyl, alkenyl or alkynyl optionally may be substituted;

n is 0-2;

each R^{2'} independently is H; optionally substituted alkyl, alkenyl, or alkynyl; -C(O)O(alkyl), -C(O)O(lower alkyl), -C(O)O(alkenyl), -C(O)O(alkynyl),
20 -C(O)NH₂, -C(O)NH(alkyl), -C(O)N(alkyl)₂, -OH, -O(acyl), -O(lower acyl), -O(alkyl), -O(lower alkyl), -O(alkenyl), halogen, halogenated alkyl and particularly CF₃, azido, cyano, NO₂, -S(alkyl), -S(alkenyl), -S(alkynyl), NH₂, -NH(alkyl), -N(alkyl)₂, -NH(alkenyl), -NH(alkynyl), -NH(acyl), or -N(acyl)₂; and

Base is selected from the group consisting of:



5

wherein

each R' and R'' independently is H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, halogen, halogenated alkyl, OH, CN, N₃, carboxy, C₁₋₄alkoxycarbonyl, NH₂, C₁₋₄ alkylamino, di(C₁₋₄ alkyl)amino, C₁₋₆ alkoxy, C₁₋₆ alkylsulfonyl, (C₁₋₄ alkyl)₀₋₂ aminomethyl;

10


each W is Cl, Br, I, F, halogenated alkyl, alkoxy, OH, SH, O-alkyl, S-alkyl, O-alkenyl, O-alkynyl, S-alkenyl, S-alkynyl, -OC(O)NR⁴R⁴, O-acyl, S-acyl, CN, SCN, OCN, NO₂, N₃, NH₂, NH(alkyl), N(alkyl)₂, NH-cycloalkyl, NH-acyl, NH=NH, CONH₂, CONH(alkyl), or CON(alkyl)₂; and

each R⁴ is independently H, acyl, or C₁₋₆ alkyl;

15

each Z is O, S, NH, N-OH, N-NH₂, NH(alkyl), N(alkyl)₂, N-cycloalkyl, alkoxy, CN, SCN, OCN, SH, NO₂, NH₂, N₃, NH=NH, NH(alkyl), N(alkyl)₂, CONH₂, CONH(alkyl), or CON(alkyl)₂.

38. The compound of claim 37, wherein $R^{2'}$ is an optionally substituted alkyl, alkenyl, or alkynyl; halogen, halogenated alkyl, CH_3 , CF_3 , azido, or cyano.
39. The compound of claim 37, wherein $R^{2'}$ is an optionally substituted alkyl, alkenyl, or alkynyl; halogen, halogenated alkyl, CH_3 , or CF_3 .
- 5 40. The compound of claim 37, wherein $R^{2'}$ is CH_3 or CF_3 .
41. The compound of claim 37, wherein each R, R^{2*} , and R^{3*} is independently H, mono-, di-, or triphosphate, a stabilized phosphate, or phosphonate.
42. The compound of claim 37, wherein each R, R^{2*} , and R^{3*} is independently H.
43. The compound of claim 37, wherein each R, R^{2*} , and R^{3*} is independently H,
10 acyl, or an amino acid acyl residue.
44. The compound of claim 37, wherein X is O or S.
45. The compound of claim 37, wherein X is O.
46. A compound of the general structure of Formula (IV):



(IV)

or a pharmacologically acceptable salt or prodrug thereof, wherein:

each R, R^{2*} , and R^{3*} independently is H, mono-, di-, or triphosphate, a stabilized phosphate, or phosphonate; optionally substituted alkyl, lower alkyl, optionally substituted alkenyl or alkynyl, acyl, -C(O)-(alkyl), -C(O)(lower alkyl), -C(O)-(alkenyl), -C(O)-(alkynyl), lipid, phospholipid, carbohydrate, peptide, cholesterol, an amino acid residue or derivative, or other pharmaceutically acceptable leaving group that is capable of providing H or phosphate when administered *in vivo*;

X is O, S[O]_n, CH₂, CHOH, CH-alkyl, CH-alkenyl, CH-alkynyl, C-dialkyl, CH-O-alkyl, CH-O-alkenyl, CH-O-alkynyl, CH-S-alkyl, CH-S-alkenyl, CH-S-alkynyl, NH, N-alkyl, N-alkenyl, N-alkynyl, S(O)N-alkyl, S(O)N-alkenyl, S(O)N-alkynyl, SCH-halogen, or C-(halogen)₂, wherein alkyl, alkenyl or alkynyl optionally may be substituted;


5

n is 0-2;


each R^{3'} independently is H; optionally substituted alkyl, alkenyl, or alkynyl; -C(O)O(alkyl), -C(O)O(lower alkyl), -C(O)O(alkenyl), -C(O)O(alkynyl), -C(O)NH₂, -C(O)NH(alkyl), -C(O)N(alkyl)₂, -OH, -O(acyl), -O(lower acyl), -O(alkyl), -O(lower alkyl), -O(alkenyl), halogen, halogenated alkyl and particularly CF₃, azido, cyano, NO₂, -S(alkyl), -S(alkenyl), -S(alkynyl), NH₂, -NH(alkyl), -N(alkyl)₂, -NH(alkenyl), -NH(alkynyl), -NH(acyl), or -N(acyl)₂; and

10


Base is selected from the group consisting of:




(i)




(ii)




(iii)




(iv)




(v)




(vi)




(vii)




(viii)



(ix)



(x)




(xi)

wherein

- each R' and R'' independently is H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, halogen, halogenated alkyl, OH, CN, N₃, carboxy, C₁₋₄alkoxycarbonyl, NH₂, C₁₋₄ alkylamino, di(C₁₋₄ alkyl)amino, C₁₋₆ alkoxy, C₁₋₆ alkylsulfonyl, (C₁₋₄ alkyl)₀₋₂ aminomethyl;
- 5 each W is Cl, Br, I, F, halogenated alkyl, alkoxy, OH, SH, O-alkyl, S-alkyl, O-alkenyl, O-alkynyl, S-alkenyl, S-alkynyl, -OC(O)NR⁴R⁴, O-acyl, S-acyl, CN, SCN, OCN, NO₂, N₃, NH₂, NH(alkyl), N(alkyl)₂, NH-cycloalkyl, NH-acyl, NH=NH, CONH₂, CONH(alkyl), or CON(alkyl)₂; and
- each R⁴ is independently H, acyl, or C₁₋₆ alkyl;
- 10 each Z is O, S, NH, N-OH, N-NH₂, NH(alkyl), N(alkyl)₂, N-cycloalkyl, alkoxy, CN, SCN, OCN, SH, NO₂, NH₂, N₃, NH=NH, NH(alkyl), N(alkyl)₂, CONH₂, CONH(alkyl), or CON(alkyl)₂.
47. The compound of claim 46, wherein R^{3'} is an optionally substituted alkyl, alkenyl, or alkynyl; halogen, halogenated alkyl, CH₃, CF₃, azido, or cyano.
- 15 48. The compound of claim 46, wherein R^{3'} is an optionally substituted alkyl, alkenyl, or alkynyl; halogen, halogenated alkyl, CH₃, or CF₃.
49. The compound of claim 46, wherein R^{3'} is CH₃ or CF₃.
50. The compound of claim 46, wherein each R, R^{2*}, and R^{3*} is independently H, mono-, di-, or triphosphate, a stabilized phosphate, or phosphonate.
- 20 51. The compound of claim 46, wherein each R, R^{2*}, and R^{3*} is independently H.
52. The compound of claim 46, wherein each R, R^{2*}, and R^{3*} is independently H, acyl, or an amino acid acyl residue.
53. The compound of claim 46, wherein X is O or S.
54. The compound of claim 46, wherein X is O.
- 25 55. A pharmaceutical composition comprising an anti-virally effective amount of a compound of any one of claims 35-37 and 46, optionally with a pharmaceutically acceptable carrier, diluent or excipient.


56. The pharmaceutical composition of claim 55 wherein the compound, salt or prodrug thereof is in the form of a dosage unit.
57. The pharmaceutical composition of claim 56 wherein the dosage unit contains from about 0.01 to about 50 mg of the compound.
- 5 58. The pharmaceutical composition of claim 57, wherein said dosage unit is a tablet or capsule.
59. The pharmaceutical composition of claim 55, further comprising one or more additional anti-virally effective agents.
- 10 60. The pharmaceutical composition of claim 59, wherein the additional anti-virally agent is selected from the group consisting of an interferon, ribavirin, an interleukin, an NS3 protease inhibitor, a cysteine protease inhibitor, a thiazolidine derivative, a thiazolidine and a benzanilide, phenanthrenequinone, a helicase inhibitor, a polymerase inhibitor, a nucleotide analogue, gliotoxin, cerulenin, an antisense oligodeoxynucleotide, an inhibitor of IRES-dependent translation, and a
15 ribozyme.
61. The pharmaceutical composition of claim 60 wherein the additional anti-virally effective agent is an interferon.
62. The pharmaceutical composition of claim 61, wherein the additional anti-virally effective agent is selected from the group consisting of pegylated interferon alpha
20 2a, interferon alphacon-1, natural interferon, albuferon, interferon beta-1a, omega interferon, interferon alpha, interferon gamma, interferon tau, interferon delta and interferon gamma- 1b.
63. The pharmaceutical composition of any one of claims 35-37 and 46, wherein the compound is in substantially pure form.
- 25 64. The pharmaceutical composition of claim 63, wherein the compound is at least 90% by weight of the β -D-isomer.
65. The pharmaceutical composition of claim 63, wherein the compound is at least 95% by weight of the β -D-isomer.

66. The pharmaceutical composition of claim 63 wherein the compound is at least 90% by weight of the β -L-isomer.
67. The pharmaceutical composition of claim 63 wherein the compound is at least 95% by weight of the β -L-isomer.
- 5 68. Use of a compound of any one of claims 35-54 for the treatment of treatment of a host infected with a *Flaviviridae*.
69. Use of a compound of any one of claims 35-54 for the manufacture of a medicament for the treatment of a host infected with a *Flaviviridae*.
- 10 70. Use of a composition of any one of claims 55-67 for the treatment of treatment of a host infected with a *Flaviviridae*.
71. Use of a composition of any one of claims 55-67 for the manufacture of a medicament for the treatment of a host infected with a *Flaviviridae*.
72. The use of any one of claims 68-71, wherein the host is a mammal.
73. The use of claim 72, wherein the mammal is a human.



(I)

or



(II)

FIGURE 1




FIGURE 2




FIGURE 3